

**PATENT APPLICATION**

**PEPTIDE EXTENDED GLYCOSYLATED POLYPEPTIDES**

**Inventor(s):**

Jens Okkels, a citizen of  
Denmark, residing at:  
147 Ellesvinget, DK-2950 Vedbaek, Denmark

Anne Dam Jensen, a citizen of  
Denmark, residing at:  
13 Svanevej, DK-2400 Copenhagen, Denmark

Bart van den Hazel, a citizen of  
Netherlands, residing at:  
29 Olufsvej, DK-2100 Copenhagen, Denmark

**Assignee:** Maxygen ApS  
1, Agern Alle, DK-2970 Hoersholm, Denmark

**Entity:** Large

**THE LAW OFFICES OF JONATHAN ALAN QUINE**

P.O. Box 458  
Alameda, CA 94501  
Internet address: [www.quinelaw.com](http://www.quinelaw.com)

Phone: (510) 337-7871  
Fax: (510) 337-7877  
E-mail: [jaquine@quinelaw.com](mailto:jaquine@quinelaw.com)

**PEPTIDE EXTENDED GLYCOSYLATED POLYPEPTIDES**

**CROSS-REFERENCES TO RELATED APPLICATIONS**

5 This application claims priority to and benefit of the following United States  
Provisional and International Patent Applications: Danish Patent Application PA 2000 01027,  
filed June 30, 2000; United States Provisional Application 60/217,497, filed July 11, 2000; PCT  
Application PCT/DK00/00743, filed December 29, 2000; PCT Application PCT/DK01/00090,  
filed February 09, 2001; Danish Patent Application PA 2000 01092, filed 14 July 2000; and  
10 United States Provisional Application 60/225,558, filed August 16, 2000, the specifications of  
which are each incorporated in their entirety for all purposes.

**FIELD OF THE INVENTION**

The present invention relates to novel glycosylated polypeptides as well as means  
and methods for their preparation.

**BACKGROUND OF THE INVENTION**

Polypeptides, including proteins, are used for a wide range of applications,  
including industrial uses and human or veterinary therapy.

One generally recognized drawback associated with many polypeptides is that  
they do not have a sufficiently high stability, are immunogenic or allergenic, have a reduced  
20 serum half-life, are susceptible to clearance, are susceptible to proteolytic degradation, and the  
like.

One method for improving properties of polypeptides has been to attach non-  
peptide moieties to the polypeptide to improve properties thereof. For instance, polymer  
molecules such as PEG has been used for reducing immunogenicity and/or increasing serum  
25 half-life of therapeutic polypeptides and for reducing allergenicity of industrial enzymes.  
Glycosylation has been suggested as another convenient route for improving properties of  
polypeptides such as stability, half-life, etc.

Machamer and Rose (1988) *J. Biol. Chem.* 263: 5948-5954 and 5955-5960,  
disclose modified glycoprotein G of vesicular stomatitis virus that is glycosylated at additional  
30 N-glycosylation sites introduced in the polypeptide backbone.

US 5,218,092 discloses physiologically active polypeptides with at least one new or additional carbohydrate attached thereto. The additional carbohydrate molecule(s) is/are provided by adding one or more additional N-glycosylation sites to the polypeptide backbone, and expressing the polypeptide in a glycosylating host cell.

5 US 5,041,376 discloses a method of identifying or shielding epitopes of a transportable protein, in which method an N-glycosylation site is introduced on the exposed surface of the protein backbone (using oligonucleotide-directed mutagenesis of the nucleotide sequence encoding the protein), the resulting protein is expressed, glycosylated and assayed for protein activity and for shielded epitopes.

10 WO 00/26354 discloses a method of reducing the allergenicity of proteins by including an additional glycosylation site in the protein backbone and glycosylating the resulting protein variant.

Guan et al. (1985) *Cell* 42: 489-496 disclose glycosylated fusion protein variants comprising a rat growth hormone backbone C-terminally extended with transmembrane and cytoplasmic domains of the vesicular stomatitis virus glycoprotein, which growth hormone backbone has been modified to incorporate two additional N-glycosylation sites.

WO 97/04079 discloses lipolytic enzymes modified to by an N- or C-terminal peptide extension capable of conferring improved performance, in particular wash performance to the enzyme.

20 Matsuura et al. (1999) *Nature Biotechnology* 17: 58-61 disclose the use of random elongation mutagenesis for improving thermostability of a non-glycosylated microbial catalase. The random elongation mutagenesis is conducted in the C-terminal end of the catalase.

US 5,338,835, entitled CTP extended forms of FSH, describe the use of the C-terminal portion of the CG beta subunit or a variant thereof for extension of the C-terminal of 25 CG, FSH and LH. Said C-terminal portion may comprise O-glycosylation sites. It is speculated that a similar approach may be used for other proteins.

US 5,508,261 discloses alpha, beta-heterodimeric polypeptide having binding affinity to vertebrate luteinizing hormone (LH) receptors and vertebrate follicle stimulating hormone (FSH) receptors comprising a glycoprotein hormone alpha-subunit polypeptide and a 30 specified non-naturally occurring beta-subunit polypeptide.

WO 95/05465 discloses EPO analogs which have one or more amino acids extending from the C-terminal end of EPO, the C-terminal extension having at least one additional carbohydrate site. The 28 amino acid C-terminal part of CG (having four O-glycosylation sites) is mentioned as an example.

5 WO 97/30161 discloses hybrid proteins comprising two coexpressed amino acid sequences forming a dimer, each comprising a) at least one amino acid sequence selected from a homomeric receptor, a chain of a heteromeric receptor, a ligand, and fragments thereof; and b) a subunit of a heterodimeric proteinaceous hormone or fragments thereof; in which a) and b) are joined directly or through a peptide linker, and, in each couple, the two subunits (b) are different  
10 and capable of aggregating to form a dimer complex.

In none of the above reference has it been disclosed or indicated that a polypeptide of interest can be modified to include additional glycosylation sites by N-terminally extending said polypeptide with a peptide sequence comprising one or more additional glycosylation sites. The present invention is based on this finding.

#### SUMMARY OF THE INVENTION

Accordingly, in a first aspect the invention relates to a glycosylated polypeptide comprising the primary structure,

$\text{NH}_2\text{-X-Pp-COOH}$

wherein

X is a peptide addition comprising or contributing to a glycosylation site, and  
Pp is a polypeptide of interest.

The introduction of additional glycosylation sites by means of a peptide addition is an elegant way of providing additional glycosylation sites in a polypeptide of interest. More specifically, the invention has the advantage that polypeptides with altered glycosylation pattern  
25 are more easily obtained, e.g., the variants can be designed without detailed knowledge or use of structural and/or functional properties of the polypeptide. Also, the utilization of glycosylation sites introduced by a peptide addition has been found to be improved relative to glycosylation sites introduced within a structural part of the polypeptide Pp. Also other properties of the peptide extended polypeptide, such as uptake in specific cells, may be improved relative to a  
30 polypeptide modified with glycosylation sites in a structural part (and not being subjected to peptide extension).

In a second aspect the invention relates to a glycosylated polypeptide comprising the primary structure  $\text{NH}_2\text{-P}_x\text{-X-P}_y\text{-COOH}$ , wherein

$\text{P}_x$  is an N-terminal part of a polypeptide Pp of interest,

$\text{P}_y$  is a C-terminal part of said polypeptide Pp, and

5 X is a peptide addition comprising or contributing to a glycosylation site.

In other aspects the invention relates to a nucleotide sequence encoding a polypeptide of the invention, an expression vector comprising said nucleotide sequence and methods of preparing a polypeptide of the invention.

10 In a further aspect the invention relates to a method of improving (a) selected property/ies of a polypeptide Pp of interest, which method comprises a) preparing a nucleotide sequence encoding a polypeptide comprising the primary structure

$\text{NH}_2\text{-X-Pp-COOH}$ ,

wherein

15 X is a peptide addition comprising or contributing to a glycosylation site, the peptide addition being capable of conferring the selected improved property/ies to the polypeptide Pp,

b) expressing the nucleotide sequence of a) in a suitable host cell under conditions ensuring attachment of an oligosaccharide moiety thereto, optionally

c) conjugating the expressed polypeptide of b) to a second non-peptide moiety,

20 and,

d) recovering the polypeptide resulting from step c).

### BRIEF DESCRIPTION OF THE FIGURES

Figure 1 is a dose response curve for uptake of glucocerebrosidase wildtype and modified according to the invention into J774E macrophages. The activity is measured by the  
25 GCB activity assay.

Figure 2 illustrates the pharmacokinetics of a FSH polypeptide produced according to the invention.

## DETAILED DISCUSSION

### DEFINITIONS

In the context of the present application and invention the following definitions apply:

5           The term “conjugate” is used to refer to the covalent attachment of one or more polypeptide(s) to one or more non-peptide moieties. The term covalent attachment means that the polypeptide and the non-peptide moiety are either directly covalently joined to one another, or else are indirectly covalently joined to one another through an intervening moiety or moieties, such as a bridge, spacer, or linkage moiety or moieties.

10           The term “non-peptide moiety” is intended to indicate a molecule, different from a peptide polymer composed of amino acid monomers and linked together by peptide bonds, which molecule is capable of conjugating to an attachment group of the polypeptide of the invention. Examples of such molecule include polymers, e.g., polyalkylene oxide moieties lipophilic groups, e.g., fatty acids and ceramides. The term “polymer molecule” is defined as a molecule formed by covalent linkage of two or more monomers and may be used interchangeably with “polymeric group.” Except where the number of non-peptide moieties, such as polymeric groups, attached to the polypeptide is expressly indicated, every reference to “non-peptide moiety” referred to herein is intended as a reference to one or more non-peptide moieties attached to the polypeptide.

20           The term “oligosaccharide moiety” is intended to indicate a carbohydrate-containing molecule comprising one or more monosaccharide residues, capable of being attached to the polypeptide (to produce a glycosylated polypeptide) by way of *in vivo* or *in vitro* glycosylation. Except where the number of oligosaccharide moieties attached to the polypeptide is expressly indicated, every reference to “oligosaccharide moiety” referred to herein is intended as a reference to one or more such moieties attached to the polypeptide.

25           The term “*in vivo* glycosylation” is intended to mean any attachment of an oligosaccharide moiety occurring *in vivo*, i.e., during posttranslational processing in a glycosylating cell used for expression of the polypeptide, e.g., by way of N-linked and O-linked glycosylation. Usually, the N-glycosylated oligosaccharide moiety has a common basic core structure composed of five monosaccharide residues, namely two N-acetylglucosamine residues and three mannose residues. The exact oligosaccharide structure depends, to a large extent, on

the glycosylating organism in question and on the specific polypeptide. Depending on the host cell in question the glycosylation is classified as a high mannose type, a complex type or a hybrid type. The term “*in vitro* glycosylation” is intended to refer to a synthetic glycosylation performed *in vitro*, normally involving covalently linking an oligosaccharide moiety to an attachment group of a polypeptide, optionally using a cross-linking agent. *In vivo* and *in vitro* glycosylation are discussed in detail further below.

An “N-glycosylation site” has the sequence N-X’-S/T/C-X’’, wherein X’ is any amino acid residue except proline, X’’ any amino acid residue that may or may not be identical to X’ and preferably is different from proline, N asparagine and S/T/C either serine, threonine or cysteine, preferably serine or threonine, and most preferably threonine. The oligosaccharide moiety is attached to the N-residue of such site. An “O-glycosylation site” is the OH-group of a serine or threonine residue. An “*in vitro* glycosylation site” is, e.g., selected from the group consisting of the N-terminal amino acid residue of the polypeptide, the C-terminal residue of the polypeptide, lysine, cysteine, arginine, glutamine, aspartic acid, glutamic acid, serine, tyrosine, histidine, phenylalanine and tryptophan. Of particular interest is an *in vitro* glycosylation site that is an epsilon-amino group, in particular as part of a lysine residue.

The term “peptide addition” is intended to indicate one or more consecutive amino acid residues that are added to the amino acid sequence of the polypeptide Pp of interest. Normally, the peptide addition is linked to the amino acid sequence of the polypeptide Pp by a peptide linkage.

The term “attachment group” is intended to indicate a functional group of the polypeptide, in particular of an amino acid residue thereof or an oligosaccharide moiety attached to the polypeptide, capable of attaching a non-peptide moiety of interest. Useful attachment groups and their matching non-peptide moieties are apparent from the table below.

The term “comprising an attachment group” is intended to mean that the attachment group is present on an amino acid residue of the relevant peptide or polypeptide or on an oligosaccharide moiety attached to said peptide or polypeptide.

The term “contributing to a glycosylation site” as used in connection with the peptide addition X is intended to cover the situation, where a glycosylation site is formed from more than one amino acid residue (as is the case with an N-glycosylation site), and where at least one such amino acid residue originates from the peptide X and at least one amino acid residue

originates from the polypeptide Pp, whereby the glycosylation site can be considered to bridge X and Pp (or, where relevant, P<sub>x</sub> or P<sub>y</sub>).

The term “non-structural part” as used about a part of the polypeptide Pp of interest is intended to indicate a part of either the C- or N-terminal end of the folded polypeptide (e.g., protein) that is outside the first structural element, such as an  $\alpha$ -helix or a  $\beta$ -sheet structure. The non-structural part can easily be identified in a three-dimensional structure or model of the polypeptide. If no structure or model is available, a non-structural part typically comprises or consists of the first or last 1-20 (1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19 or 20) amino acid residues, such as 1-10 amino acid residues of the amino acid sequence constituting the mature form of the polypeptide of interest.

Amino acid names and atom names (e.g., CA, CB, NZ, N, O, C, etc) are used as defined by the Protein DataBank (PDB) ([www.pdb.org](http://www.pdb.org)) which are based on the IUPAC nomenclature (IUPAC Nomenclature and Symbolism for Amino Acids and Peptides (residue names, atom names e.t.c.), *Eur. J. Biochem.*, **138**, 9-37 (1984) together with their corrections in *Eur. J. Biochem.*, **152**, 1 (1985). The term “amino acid residue” is intended to indicate an amino acid residue contained in the group consisting of alanine (Ala or A), cysteine (Cys or C), aspartic acid (Asp or D), glutamic acid (Glu or E), phenylalanine (Phe or F), glycine (Gly or G), histidine (His or H), isoleucine (Ile or I), lysine (Lys or K), leucine (Leu or L), methionine (Met or M), asparagine (Asn or N), proline (Pro or P), glutamine (Gln or Q), arginine (Arg or R), serine (Ser or S), threonine (Thr or T), valine (Val or V), tryptophan (Trp or W), and tyrosine (Tyr or Y) residues. The terminology used for identifying amino acid positions/mutations is illustrated as follows: A15 (indicates an alanine residue in position 15 of the polypeptide), A15T (indicates replacement of the alanine residue in position 15 with a threonine residue), A15[T/S] (indicates replacement of the alanine residue in position 15 with a threonine residue or a serine residue). Multiple substitutions are indicated with a “+,” e.g., A15T+F57S means an amino acid sequence which comprises a substitution of the alanine residue in position 15 for a threonine residue and a substitution of the phenylalanine residue in position 57 for a serine residue.

The term “nucleotide sequence” is intended to indicate a consecutive stretch of two or more nucleotides. The nucleotide sequence can be of genomic, cDNA, RNA, semisynthetic, synthetic origin, or any combinations thereof.

“Cell,” “host cell,” “cell line” and “cell culture” are used interchangeably herein and all such terms should be understood to include progeny resulting from growth or culturing of a cell. “Transformation” and “transfection” are used interchangeably to refer to the process of introducing DNA into a cell.

5 “Operably linked” refers to the covalent joining of two or more nucleotide sequences in such a manner that the normal function of the sequences can be performed. For example, the nucleotide sequence encoding a presequence or secretory leader is operably linked to a nucleotide sequence for a polypeptide if it is expressed as a preprotein that participates in the secretion of the polypeptide: a promoter or enhancer is operably linked to a coding sequence if it  
10 affects the transcription of the sequence.

“Introduction” or “removal” of a glycosylation site or an attachment group for a non-peptide moiety is normally achieved by introducing or removing an amino acid residue comprising or contributing to such site or group to/from the relevant amino acid sequence, conveniently by suitable modification of the encoding nucleotide sequence. For instance, when an N-glycosylation site is to be introduced/removed this can be done by introducing/removing a codon for the amino acid residue(s) required for a functional N-glycosylation site. When an attachment group for a PEG molecule is to be introduced/removed, it will be understood that this  
15 be done by introducing/removing a codon for an amino acid residue, e.g., a lysine residue, comprising such group to/from the encoding nucleotide sequence. The term “introduce” is primarily intended to include substitution of an existing amino acid residue, but can also mean insertion of additional amino acid residue. The term “remove” is primarily intended to include substitution of the amino acid residue to be removed for another amino acid residue, but can also mean deletion (without substitution) of the amino acid residue to be removed.

The term “epitope” is used in its conventional meaning to indicate one or more  
25 amino acid residue(s) displaying specific 3D and/or charge characteristics at the surface of the polypeptide, which is/are capable of giving rise to an immune response in a mammal and/or specifically binding to an antibody raised against said epitope or which is/are capable of giving rise to an allergic response.

The term “unshielded epitope” is intended to indicate that the epitope is not  
30 shielded and therefore has the above properties. The term “shielded epitope” is intended to indicate that the non-peptide moiety shields, and thus inactivates the epitope, whereby it is no

longer capable of giving rise to any substantial immune response in a mammal, e.g., due to inappropriate processing and/or presentation in the antigen presenting cells, and/or of reacting with an antibody raised against the unshielded epitope. The shielding should thus be effective in both the naïve mammal and mammals that already produce antibodies reacting with the unshielded epitope.

The degree of shielding of epitopes can be determined as reduced immunogenicity and/or reduced antibody reactivity and/or reduced reactivity with monoclonal antibodies raised against the epitope(s) in question using methods known in the art. The degree of shielding of allergenic epitopes can be determined, e.g., as described in WO 00/26354.

The term “reduced” as used about an immunogenic or allergic response is intended to indicate that a given molecule gives rise to a measurably lower immune or allergic response than a reference molecule, when determined under comparable conditions. Preferably, the relevant response is reduced by at least 25%, such as at least 50%, such as preferably by at least 75%, such as by at least 90% or even at least 100%.

The term “serum half-life” is used in its normal meaning, i.e., the time in which half of the relevant molecules circulate in the plasma or bloodstream prior to being cleared. Alternatively used terms include “plasma half-life,” “circulating half-life,” “serum clearance,” “plasma clearance” and “clearance half-life.” The term “functional *in vivo* half-life” is the time in which 50% of a given function (such as biological activity) of the relevant molecule is retained, when tested *in vivo* (such as the time at which 50% of the biological activity of the molecule is still present in the body/target organ, or the time at which the activity of the polypeptide is 50% of the initial value). The molecule is normally cleared by the action of one or more of the reticuloendothelial systems (RES), kidney (e.g., by glomerular filtration), spleen or liver, or receptor-mediated elimination, or degraded by specific or unspecific proteolysis.

Normally, clearance depends on size or hydrodynamic volume (relative to the cut-off for glomerular filtration), shape/rigidity, charge, attached carbohydrate chains, and the presence of cellular receptors for the molecule. The term “increased” as used about serum half-life or functional *in vivo* half-life is used to indicate that the relevant half-life of the relevant molecule is statistically significantly increased relative to that of the reference molecule as determined under comparable conditions. For instance, the relevant half-life is increased by at least 25%, such as by at least 50%, by at least 100% or by at least 1000%.

The term “function” is intended to indicate one or more specific functions of the polypeptide of interest and is to be understood qualitatively (i.e., having a similar function as the polypeptide of interest) and not necessarily quantitatively (i.e., the magnitude of the function is not necessarily similar). Typically, a given polypeptide has many different functions, examples of which are given further below in the section entitled “Screening for or measurement of function.” For therapeutically useful polypeptides an important “function” is biological activity, e.g., *in vitro* or *in vivo* bioactivity. For enzymes, an important function is biological activity such as catalytic activity.

The interchangeably used terms “measurable function” and “functional” are intended to indicate that the relevant function (preferably reflecting the intended use) of a polypeptide of the invention is above detection limit when measured by standard methods known in the art, e.g., as an *in vitro* bioactivity and/or *in vivo* bioactivity. For instance, if the polypeptide is a hormone and the function of interest is the hormone’s affinity towards a specific receptor a measurable function is defined to be a detectable affinity between the hormone modified in accordance with the invention and the receptor as determined by the normal methods used for measuring such affinity. If the polypeptide is an enzyme and a function of interest is the catalytic activity a measurable function is the enzyme’s ability to catalyze a reaction involving the normal substrates for the enzyme as measured by the normal methods for determining the enzyme activity in question. Typically, if not otherwise stated herein, a measurable function is at least 2%, such as at least 5% of that of the unmodified polypeptide Pp, as determined under comparable conditions, e.g., in the range of 2-1000%, such as 2-500% or 2-100%, such as 5-100% of that of the unmodified polypeptide.

The term “functional site” is intended to indicate one or more amino acid residues which is/are essential for or otherwise involved in the function or performance of the polypeptide, i.e., the amino acid residue(s) that mediate(s) a desired biological activity of the polypeptide Pp. Such amino acid residues are “located at” the functional site. For instance, the functional site can be a binding site (e.g., a receptor-binding site of a hormone or growth factor or a ligand-binding site of a receptor), a catalytic site (e.g., of an enzyme), an antigen-binding site (e.g., of an antibody), a regulatory site (e.g., of a polypeptide subject to regulation), or an interaction site (e.g., for a regulatory protein or an inhibitor). The functional site can be

determined by methods known in the art and is conveniently identified by analysing a three-dimensional or model structure of the polypeptide complexed to a relevant ligand.

The term “polypeptide” is intended to indicate any structural form (e.g., the primary, secondary or tertiary form (i.e., protein form)) of an amino acid sequence comprising more than 5 amino acid residues, which may or may not be post-translationally modified (e.g., acetylated, carboxylated, phosphorylated, lipidated, or acylated). The interchangeably used terms “native” and “wild-type” are used about a polypeptide which has an amino acid sequence that is identical to one found in nature. The native polypeptide is typically isolated from a naturally occurring source, in particular a mammalian or microbial source, such as a human source, or is produced recombinantly by use of a nucleotide sequence encoding the naturally occurring amino acid sequence. The term “native” is intended to encompass allelic variants of the polypeptide in question. A “variant” is a polypeptide, which has an amino acid sequence that differs from that of a native polypeptide in one or more amino acid residues. The variant is typically prepared by modification of a nucleotide sequence encoding the native polypeptide (e.g., to result in substitution, deletion or truncation of one or more amino acid residues of the polypeptide or by introduction (by addition or insertion) of one or more amino acid residues into the polypeptide) so as to modify the amino acid sequence constituting said native polypeptide. A “fragment” is a part of a parent native or variant polypeptide, typically differing from such parent in one or more removed C-terminal or N-terminal amino acid residues or removal of both types of such residues. Normally, the variant or fragment has retained at least one of the functions of the corresponding parent polypeptide (e.g., a biological function such as enzyme activity or receptor binding capability). Normally, the polypeptide Pp is a full length protein or a variant or fragment thereof.

The term “antibody” includes single monoclonal antibodies (including agonist and antagonist antibodies) and antibody compositions with polyepitopic specificity (also termed polyclonal antibodies).

The term “monoclonal antibody” is used in its conventional meaning to indicate a population of substantially homogeneous antibodies. The individual antibodies comprised in the population have identical binding affinities and vary structurally only to a limited extent. Monoclonal antibodies are highly specific, being directed against a single epitope. Furthermore, in contrast to conventional (polyclonal) antibody preparations that typically include different

antibodies directed against different epitopes, each monoclonal antibody is directed against a single epitope on the antigen. The antibody to be modified is preferably a human or humanized monoclonal antibody.

“Antibody fragment” is defined as a portion of an intact antibody comprising the antigen binding site or the entire or part of the variable region of the intact antibody, wherein the portion is free of the constant heavy chain domains (i.e., CH2, CH3, and CH4, depending on antibody isotype) of the Fc regions of the intact antibody. Examples of antibody fragments include Fab, Fab', Fab'-SH, F(ab')2, and Fv fragments; diabodies; any antibody fragment that is a polypeptide having a primary structure consisting of one uninterrupted sequence of contiguous amino acid residues (which may also be termed a single chain antibody fragment or a single chain polypeptide).

### POLYPEPTIDE OF THE INVENTION

In its first aspect the invention relates to a glycosylated polypeptide comprising the primary structure:

$\text{NH}_2\text{-X-Pp-COOH}$ ,

wherein X is a peptide addition comprising or contributing to a glycosylation site, and Pp is a polypeptide of interest.

In one embodiment, the polypeptide consists essentially of or consists of a polypeptide with the primary structure  $\text{NH}_2\text{-X-Pp-COOH}$ .

The peptide addition according to this aspect is preferably one, which has less than 90% identity to a native full length protein. The identity is determined on the basis of an alignment of the peptide addition to the entire amino acid sequence of the full length native protein, the alignment being made to ensure the highest possible degree of identity between amino acid residues. For instance, the program CLUSTALW version 1.74 using default parameters (Thompson et al. (1994) CLUSTAL W: improving the sensitivity of progressive multiple sequence alignment through sequence weighting, position-specific gap penalties and weight matrix choice *Nucleic Acids Research* 22:4673-4680) can be used.

Usually, the peptide addition is fused to the N-terminal end of the polypeptide Pp as reflected in the above shown structure so as to provide an N-terminal elongation of the polypeptide Pp. However, it is also possible to insert the peptide addition within the amino acid sequence of the polypeptide Pp. This is reflected in the polypeptide according to the second

aspect of the invention, wherein the polypeptide comprises the primary structure  $\text{NH}_2\text{-P}_x\text{-X-P}_y\text{-COOH}$ , wherein

$\text{P}_x$  is an N-terminal part of a polypeptide Pp of interest,

$\text{P}_y$  is a C-terminal part of said polypeptide Pp, and

5 X is a peptide addition comprising or contributing to a glycosylation site.

In one embodiment, the polypeptide consists essentially of or consists of a polypeptide with the primary structure  $\text{NH}_2\text{-P}_x\text{-X-P}_y\text{-COOH}$ .

In order to minimize structural changes effected by the insertion of the peptide addition within the sequence of the polypeptide Pp, it is desirable that it be inserted in a non-  
10 structural part thereof. For instance,  $\text{P}_x$  is a non-structural N-terminal part of a mature polypeptide Pp, and  $\text{P}_y$  is a structural C-terminal part of said mature polypeptide, or  $\text{P}_x$  is a structural N-terminal part of a mature polypeptide Pp, and  $\text{P}_y$  is a non-structural C-terminal part of said mature polypeptide. Preferably, when the glycosylation site to be introduced is an N-glycosylation site,  $\text{P}_x$  is a non-structural N-terminal part since, in general, the best N-glycosylation is obtained in the N-terminal part of a polypeptide.

When the peptide addition comprises only few amino acid residues, e.g., 1-5 such as 1-3 amino acid residues, and in particular 1 amino acid residue, the peptide addition can be inserted into a loop structure of the polypeptide Pp and thereby elongate said loop. When the peptide addition is constituted by one amino acid residue it will be understood that this is  
20 selected so as to ensure that a functional glycosylation site is introduced.

Polypeptides of the invention are glycosylated polypeptides. Normally, the peptide addition part of the polypeptide of the invention has attached at least one oligosaccharide moiety. The polypeptide Pp part of the polypeptide may or may not have attached at least one oligosaccharide moiety. Glycosylation can be achieved as described in the section entitled  
25 "Glycosylation"

Preferably, the polypeptide of the invention has properties such as size, charge, molecular weight and/or hydrodynamic volume that are sufficient to reduce or escape clearance by any of the clearance mechanisms disclosed herein, in particular renal clearance. Such properties are, e.g., determinable by the nature and number of oligosaccharide and second non-  
30 peptide moieties attached thereto. In one embodiment, the polypeptide of the invention has a molecular weight of at least 67 kDa, in particular at least 70 kDa as measured by SDS-PAGE

according to Laemmli, U.K., Nature Vol 227 (1970), p680-85. This is of particular relevance when the polypeptide of interest is a therapeutically useful protein, the functional *in vivo* half-life of which is to be prolonged. A molecular weight of at least 67 kDa is obtainable by introduction of a sufficient number of glycosylation sites to obtain a glycosylated polypeptide with such MW, or by conjugating the glycosylated polypeptide to a sufficient number and type of a second non-peptide moiety to obtain such MW. For instance, for a glycosylated polypeptide of interest having a molecular weight of at least 25 kDa linked to a peptide addition of 2 kDa, the combined extended polypeptide having at least two PEG-attachment groups, conjugation to two or more PEG molecules each having a molecular weight of 20 kDa results in a total molecular weight of at least 67 kDa.

Preferably, the polypeptide of the invention has at least one of the following properties relative to the polypeptide Pp, the properties being measured under comparable conditions: *in vitro* bioactivity which is at least 25%, such as at least 30% or at least 45% of that of the polypeptide Pp as measured under comparable conditions, increased affinity for a mannose receptor, a mannose-6-phosphate receptor or other carbohydrate receptors, increased serum half-life, increased functional *in vivo* half-life, reduced renal clearance, reduced immunogenicity, increased resistance to proteolytic cleavage, improved targeting to lysosomes, macrophages and/or other subpopulations of human cells, improved stability in production, improved shelf life, improved formulation, e.g., liquid formulation, improved purification, improved solubility, and/or improved expression.

Improved properties are determined by conventional methods known in the art for determining such properties. The improvement is of a magnitude that is within detection limits.

Improved affinity for or uptake by the mannose receptor is expected to result in increased uptake in phagocytic cells, preferably monocytes, macrophages (e.g., Kupffer cells, glia/microglia, alveolar phagocytes, reticulum cells, or other peripheral macrophages) or macrophage like cells (for instance osteoclasts, dendritic cells, or astrocytes) in increased uptake of the polypeptide in phagocytic cells (e.g., macrophages). This is of particular relevance when the polypeptide of interest is one for which such uptake is required for the polypeptide to exert its biological activity. Such polypeptide is e.g., an antigen intended for use for vaccine purposes or a lysosomal enzyme.

### Polypeptide of interest

The present invention can be applied broadly. Thus, the polypeptide of interest can have any function and be of any origin. Accordingly, the polypeptide can be a protein, in particular a mature protein or a precursor form thereof or a functional fragment thereof that essentially has retained a biological activity of the mature protein. Furthermore, the polypeptide can be an oligopeptide that contains in the range of 30 to 4500 amino acids, preferably in the range of 40 to 3000 amino acids.

The polypeptide can be a native polypeptide or a variant thereof. For instance, the polypeptide is a variant that comprises at least one introduced and/or at least one removed glycosylation site as compared to the corresponding native polypeptide. The variant has retained at least one function of the corresponding native polypeptide, in particular a biological activity thereof.

The polypeptide can be a therapeutic polypeptide useful in human or veterinary therapy, i.e., a polypeptide that is physiologically active when introduced into the circulatory system of or otherwise administered to a human or an animal; a diagnostic polypeptide useful in diagnosis; or an industrial polypeptide useful for industrial purposes, such as in the manufacture of goods wherein the polypeptide constitutes a functional ingredient or wherein the polypeptide is used for processing or other modification of raw ingredients during the manufacturing process.

The polypeptide can be of mammalian origin, e.g., of human, porcine, ovine, urcine, murine, rabbit, donkey, or bat origin, of microbial origin, e.g., of fungal, yeast or bacterial origin, or can be derived from other sources such as venom, leech, frog or mosquito origin. Preferably, the industrial polypeptide of interest is of microbial origin and the therapeutic polypeptide of human origin.

Specific examples of groups of polypeptides to be modified according to the invention include: an antibody or antibody fragment, an immunoglobulin or immunoglobulin fragment, a plasma protein, an erythrocyte or thrombocyte protein, a cytokine, a growth factor, a profibrinolytic protein, a binding protein, a protease inhibitor, an antigen, an enzyme, a ligand, a receptor, or a hormone. Of particular interest is a polypeptide that mediates its biological effect by binding to a cellular receptor, when administered to a patient. The antibody can be a polyclonal or monoclonal antibody, and can be of any origin including human, rabbit and murine origin. Preferably, the antibody is a human or humanized monoclonal antibody.

Immunoglobulins of interest include IgG, IgE, IgM, IgA, and IgD and fragments thereof, e.g., Fab fragments. Specific antibodies and fragments thereof are those reactive with any of the proteins mentioned immediately below.

The non-antibody polypeptide of interest can be i) a plasma protein, e.g., a factor from the coagulation system, such as Factor VII, Factor VIII, Factor IX, Factor X, Factor XIII, thrombin, protein C, antithrombin III or heparin co-factor II, Tissue factor inhibitor (e.g., 1 or 2), endothelial cell surface protein C receptor, a factor from the fibrinolytic system such as pro-urokinase, urokinase, tissue plasminogen activator, plasminogen activator inhibitor 1 (PAI-1) or plasminogen activator inhibitor 2 (PAI-2), the Von Willebrand factor, or an  $\alpha$ -1-proteinase inhibitor, ii) a erythrocyte or thrombocyte protein, e.g., hemoglobin, thrombospondin or platelet factor 4, iii) a cytokine, e.g., an interleukin such as IL-1 (e.g., IL-1 $\alpha$  or IL-1 $\beta$ ), IL-2, IL-4, IL-5, IL-6, IL-9, IL-10, IL-11, IL-12, IL-13, IL-15, IL-16, IL-17, IL-18, IL-19, IL-20, IL-21, IL-22, IL-23, a cytokine-related polypeptide, such as IL-1Ra, an interferon such as interferon- $\alpha$ , interferon- $\beta$  or interferon- $\gamma$ , a colony-stimulating factor such as GM-CSF or G-CSF, stem cell factor (SCF), a binding protein, a member of the tumor necrosis factor family (e.g TNF- $\alpha$ , lymphotoxin- $\alpha$ , lymphotoxin- $\beta$ , FasL, CD40L, CD30L, CD27L, Ox40L, 4-1BBL, RANKL, TRAIL, TWEAK, LIGHT, TRANCE, APRIL, THANK or TALL-1), iv) a growth factor, e.g platelet-derived growth factor (PDGF), transforming growth factor  $\alpha$  (TGF- $\alpha$ ), transforming growth factor  $\beta$  (TGF- $\beta$ ), epidermal growth factor (EGF), vascular endothelial growth factor (VEGF), somatotropin (growth hormone), a somatomedin such as insulin-like growth factor I (IGF-I) or insulin-like growth factor II (IGF-II), erythropoietin (EPO), thrombopoietin (TPO) or angiopoietin, v) a profibrinolytic protein, e.g., staphylokinase or streptokinase, vi) a protease inhibitor, e.g., aprotinin or CI-2A, vii) an enzyme, e.g., superoxide dismutase, catalase, uricase, bilirubin oxidase, trypsin, papain, asparaginase, arginase, arginine deiminase, adenosin deaminase, ribonuclease, alkaline phosphatase,  $\beta$ -glucuronidase, purine nucleoside phosphorylase or batroxobin, viii) an opioid, e.g., endorphins, enkephalins or non-natural opioids, ix) a hormone or neuropeptide, e.g., insulin, calcitonin, glucagons, adrenocorticotrophic hormone (ACTH), somatostatin, gastrins, cholecystokinins, parathyroid hormone (PTH), luteinizing hormone (LH), follicle-stimulating hormone (FSH), gonadotropin-releasing hormone, chorionic gonadotropin, corticotropin-releasing factor, vasopressin, oxytocin, antidiuretic hormones, thyroid-stimulating hormone, thyrotropin-releasing hormone, relaxin, glucagon-like

peptide 1 (GLP-1), glucagon-like peptide 2 (GLP-2), prolactin, neuropeptide Y, peptide YY, pancreatic polypeptide, leptin, orexin, CART (cocaine and amphetamine regulated transcript), a CART-related peptide, melanocortins (melanocyte-stimulating hormones), melanin-concentrating hormone, natriuretic peptides, adrenomedullin, endothelin, exendin, secretin, amylin (IAPP; islet amyloid polypeptide precursor), vasoactive intestinal peptide (VIP), pituitary adenylate cyclase activating polypeptide (PACAP), agouti and agouti-related peptides or somatotropin-releasing hormones, or x) another type of protein or peptide such as thymosin, bombesin, bombesin-like peptides, heparin-binding protein, soluble CD4, pigmentary hormones, hypothalamic releasing factor, melanotons, phospholipase activating protein, a detoxifying enzyme such as acylglyoxal hydrolase, or an antimicrobial peptide.

One group of polypeptides of particular interest in the present invention is selected from the group of lysosomal enzymes (as defined in US 5,929,304) such as those responsible for or otherwise involved in a lysosomal storage disease, i.e., enzymes that have a therapeutic effect on patients with a lysosomal storage disease. Such enzymes, e.g., include glucocerebrosidase,  $\alpha$ -L-iduronidase, acid  $\alpha$ -glucosidase,  $\alpha$ -galactosidase, acid sphingomyelinase, galactocerebrosidase, arylsulphatase A, sialidase, and hexosaminidase. Also, other proteins involved in lysosomal storage diseases such as Saposin A, B, C or D (Nakano et al., J. Biochem. (Tokyo) 105, 152-154, 1989; Gavrieli-Rorman and Grabowski, Genomics 5, 486-492, 1989) can be modified as described herein. Preferably, these polypeptides are of human origin.

The present inventors have shown that providing such enzymes with additional N-linked oligosaccharide moieties considerably improve properties thereof, such as stability, targeting, expression, and *in vivo* activity and targeting. Accordingly, in one embodiment, the polypeptide of the invention is a glycosylated lysosomal enzyme comprising a peptide addition comprising or contributing to a glycosylation site.

The industrial polypeptide is typically an enzyme, in particular a microbial enzyme, and can be used in products or in the manufacture of products such as detergents, household articles, personal care products, agrochemicals, textile, food products, in particular bakery products, feed products, or in industrial processes such as hard surface cleaning. The industrial polypeptide is normally not intended for internal administration to humans or animals. Specific examples include hydrolases, such as proteases, lipases or cutinases, oxidoreductases,

such as laccase and peroxidase, transferases such as transglutaminases, isomerases, such as protein disulphide isomerase and glucose isomerase, cell wall degrading enzymes such as cellulases, xylanases, pectinases, mannanases, etc., amylolytic enzymes such as endoamylases, e.g., alpha-amylases, or exo-amylases, e.g., beta-amylases or amyloglucosidases, etc. Further specific examples are those listed in WO 00/26354, the contents of which are incorporated herein by reference. Normally, an enzyme modified according to the present invention has one or more improved properties selected from the group consisting of increased stability (in particular against proteolytic degradation or thermal degradation) leading to, e.g., improved shelf life and improved performance in use; improved production, e.g., in terms of improved expression (e.g., as a consequence of improved secretion and/or increased stability of the expressed enzyme) and improved purification, decreased allergenicity, increased activity in the relevant industrial process in which it is used, and improved properties with respect to immobilization.

When the polypeptide Pp is an industrial enzyme the N-terminal peptide addition may comprise or contribute to a glycosylation site. However, it is also within the scope of the present invention to provide a polypeptide comprising an industrial enzyme and a C-terminal or N-terminal peptide addition comprising an attachment group for a second non-peptide moiety being a polymer, e.g., PEG. The peptide addition may or may not comprise a glycosylation site. The peptide addition is preferably as described herein. For instance, such attachment group can be provided by a lysine or cysteine residue.

In one embodiment, the polypeptide of the invention comprises a personal care enzyme (i.e., an enzyme useful for personal care applications), which polypeptide is incapable of passing the mucous membrane of a mammal, in particular a human exposed to the polypeptide. Thereby, allergenicity can be reduced or avoided. Furthermore, stability of such enzyme can be increased. The polypeptide according to this embodiment comprises an N-terminal or C-terminal peptide addition comprising or contributing to a glycosylation site and/or an attachment group for a second non-peptide moiety, e.g., a polymer such as PEG.

In another embodiment, the polypeptide comprises a lipase as disclosed in WO 97/04079, in particular a *Humicola lanuginosa* lipase, wherein the N- or C-terminal peptide addition comprises a glycosylation site and/or at least one attachment group for a second non-peptide moiety, e.g., a polymer such as PEG. Thereby, the N- or C-terminal peptide addition is shielded from degradation and/or increased expression, including secretion, of the enzyme is

likely to be obtained. In connection with this embodiment, the N-terminal peptide addition can comprise any of the peptide additions disclosed in WO 97/04079.

In yet another embodiment, the polypeptide Pp is an amyloglucosidase and the N- or C-terminal peptide addition comprises or contributes to a glycosylation site and/or an attachment group for a second non-peptide moiety, e.g., a polymer such as PEG. When the peptide addition is N-terminal the modification of such enzyme is contemplated to result in reduced or no degradation of the N-terminus of said enzyme (an otherwise well known problem associated with the recombinant production of amyloglucosidase). In other words, the N-terminus of the enzyme is protected by the non-peptide moiety attached to the N-terminal peptide addition of the amyloglucosidase.

In yet another embodiment, the polypeptide Pp is an antigen, in particular an antigen intended for use in eliciting an immune response (for vaccine purposes). It is contemplated to be advantageous to add N-terminal glycosylation site(s) to antigens in accordance with the invention in that the risk of changing antigenicity is thereby reduced. Antigens are recognized by a wide range of target cells, including antigen presenting cells (APC), and taken up by those cells for efficient intracellular processing and presentation to other cells of the immune system, such as, e.g., T cells, to induce or elicit desired immune responses. Antigens (and fragments thereof, e.g., antigen peptides) can be modified by a peptide addition and non-peptide moieties according to the invention. Such modifications facilitate and/or optimize uptake and/or targeting to processing compartment of the antigen by such target cells. For example, N-terminally extended antigen polypeptides of the invention are taken up by the target cells more efficiently and/or at an enhanced or improved rate (when the non-peptide moiety is one involved in such uptake). Such efficient, improved, or enhanced uptake of modified antigens by the target cells increases the kinetics and potency of the immune response to the immunizing antigen. These modifications to antigens also improve the affinity of the antigens for particular cellular receptors on target cells, including, e.g., mannose receptors and other carbohydrate receptors (in particular when the non-peptide moiety is an oligosaccharide moiety).

Antigen polypeptides of the invention include, but are not limited to those, for which an improved, enhanced or altered uptake of antigens in the following type of target cells is desired: antigen-presenting and antigen-processing cells, such as monocytes, B cells, antigen-

presenting macrophages, marginal zone macrophages, follicular dendritic cells, dendritic cells, Langerhans cells, keratinocytes, M-cells (e.g., M-cells of the gut), myocytes for intramuscular immunization or epithelial cells for mucosal immunization, Kupffer cells in the liver, and the like. A number of other cells, including capillary endothelium and some endocrine cells, can present antigen in some circumstances; the cells develop MHC class II molecules that confer antigen-presenting function. Furthermore, MHC class I molecules are expressed on the surface of most nucleated cells, including, for example, muscle cells, and therefore these cells can also present antigens to CD8+ T cells. Activated T cells, which release IFN-gamma actively induce expression of MHC molecules on some tissue cells. Such cells are also of use with the novel polypeptides of the invention. Preferably, such cells are of mammalian origin, in particular human (for use in immunization of a human) or animal (for veterinary purposes).

A wide range of antigens can be modified according to the invention. Examples are as follows:

#### Cancer antigens

Examples of cancer antigens that can be modified according to the invention include, but are not limited to: bullous pemphigoid antigen 2, prostate mucin antigen (PMA) (Beckett and Wright (1995) *Int. J. Cancer* 62: 703-710), tumor associated Thomsen-Friedenreich antigen (Dahlenborg *et al.* (1997) *Int. J. Cancer* 70: 63-71), prostate-specific antigen (PSA) (Dannull and Beldegrun (1997) *Br. J. Urol.* 1: 97-103), EpCam/KSA antigen, luminal epithelial antigen (LEA.135) of breast carcinoma and bladder transitional cell carcinoma (TCC) (Jones *et al.* (1997) *Anticancer Res.* 17: 685-687), cancer-associated serum antigen (CASA) and cancer antigen 125 (CA 125) (Kierkegaard *et al.* (1995) *Gynecol. Oncol.* 59: 251-254), the epithelial glycoprotein 40 (EGP40) (Kievit *et al.* (1997) *Int. J. Cancer* 71: 237-245), squamous cell carcinoma antigen (SCC) (Lozza *et al.* (1997) *Anticancer Res.* 17: 525-529), cathepsin E (Mota *et al.* (1997) *Am. J. Pathol.* 150: 1223-1229), tyrosinase in melanoma (Fishman *et al.* (1997) *Cancer* 79: 1461-1464), cell nuclear antigen (PCNA) of cerebral cavernomas (Notelet *et al.* (1997) *Surg. Neurol.* 47: 364-370), DF3/MUC1 breast cancer antigen (Apostolopoulos *et al.* (1996) *Immunol. Cell. Biol.* 74: 457-464; Pandey *et al.* (1995) *Cancer Res.* 55: 4000-4003), carcinoembryonic antigen (Paone *et al.* (1996) *J. Cancer Res. Clin. Oncol.* 122: 499-503; Schlom *et al.* (1996) *Breast Cancer Res. Treat.* 38: 27-39), tumor-associated antigen CA 19-9 (Tolliver and O'Brien (1997) *South Med. J.* 90: 89-90; Tsuruta *et al.* (1997) *Urol. Int.* 58: 20-24),

human melanoma antigens MART-1/Melan-A27-35 and gp100 (Kawakami and Rosenberg (1997) *Int. Rev. Immunol.* 14: 173-192; Zajac *et al.* (1997) *Int. J. Cancer* 71: 491-496), the T and Tn pancarcinoma (CA) glycopeptide epitopes (Springer (1995) *Crit. Rev. Oncog.* 6: 57-85), a 35 kD tumor-associated autoantigen in papillary thyroid carcinoma (Lucas *et al.* (1996) *Anticancer Res.* 16: 2493-2496), KH-1 adenocarcinoma antigen (Deshpande and Danishefsky (1997) *Nature* 387: 164-166), the A60 mycobacterial antigen (Maes *et al.* (1996) *J. Cancer Res. Clin. Oncol.* 122: 296-300), heat shock proteins (HSPs) (Blachere and Srivastava (1995) *Semin. Cancer Biol.* 6: 349-355), and MAGE, tyrosinase, melan-A and gp75 and mutant oncogene products (e.g., p53, ras, and HER-2/neu (Bueler and Mulligan (1996) *Mol. Med.* 2: 545-555; Lewis and Houghton (1995) *Semin. Cancer Biol.* 6: 321-327; Theobald *et al.* (1995) *Proc. Nat'l. Acad. Sci. USA* 92: 11993-11997); TAG-72, a mucin ag expressed in most human adenocarcinomas (McGuinness *et al.* (1999) *Hum Gene Ther* 10:165-73).

### Bacterial antigens

Bacterial antigens that can be modified according to the invention include, but are not limited to, *Helicobacter pylori* antigens CagA and VacA (Blaser (1996) *Aliment. Pharmacol. Ther.* 1: 73-7; Blaser and Crabtree (1996) *Am. J. Clin. Pathol.* 106: 565-7; Censini *et al.* (1996) *Proc. Nat'l. Acad. Sci. USA* 93: 14648-14643). Other suitable *H. pylori* antigens include, for example, four immunoreactive proteins of 45-65 kDa as reported by Chatha *et al.* (1997) *Indian J. Med. Res.* 105: 170-175 and the *H. pylori* GroES homologue (HspA) (Kansau *et al.* (1996) *Mol. Microbiol.* 22: 1013-1023. Other suitable bacterial antigens include, but are not limited to, the 43-kDa and the fimbriin (41 kDa) proteins of *P. gingivalis* (Boutsli *et al.* (1996) *Oral Microbiol. Immunol.* 11: 236-241); pneumococcal surface protein A (Briles *et al.* (1996) *Ann. NY Acad. Sci.* 797: 118-126); *Chlamydia psittaci* antigens, 80-90 kDa protein and 110 kDa protein (Buendia *et al.* (1997) *FEMS Microbiol. Lett.* 150: 113-9); the chlamydial exoglycolipid antigen (GLXA) (Whittum-Hudson *et al.* (1996) *Nature Med.* 2: 1116-1121); *Chlamydia pneumoniae* species-specific antigens in the molecular weight ranges 92-98, 51-55, 43-46 and 31.5-33 kDa and genus-specific antigens in the ranges 12, 26 and 65-70 kDa (Halme *et al.* (1997) *Scand. J. Immunol.* 45: 378-84); *Neisseria gonorrhoeae* (GC) or *Escherichia coli* phase-variable opacity (Opa) proteins (Chen and Gotschlich (1996) *Proc. Nat'l. Acad. Sci. USA* 93: 14851-14856), any of the twelve immunodominant proteins of *Schistosoma mansoni* (ranging in molecular weight from 14 to 208 kDa) as described by Cutts and Wilson (1997) *Parasitology* 114: 245-55; the 17-

kDa protein antigen of *Brucella abortus* (De Mot *et al.* (1996) *Curr. Microbiol.* 33: 26-30); a  
 gene homolog of the 17-kDa protein antigen of the Gram-negative pathogen *Brucella abortus*  
 identified in the nocardioform actinomycete *Rhodococcus* sp. NI86/21 (De Mot *et al.* (1996)  
*Curr. Microbiol.* 33: 26-30); the staphylococcal enterotoxins (SEs) (Wood *et al.* (1997) *FEMS*  
 5 *Immunol. Med. Microbiol.* 17: 1-10), a 42-kDa *M. hyopneumoniae* NrdF ribonucleotide  
 reductase R2 protein or 15-kDa subunit protein of *M. hyopneumoniae* (Fagan *et al.* (1997) *Infect.*  
*Immun.* 65: 2502-2507), the meningococcal antigen PorA protein (Feavers *et al.* (1997) *Clin.*  
*Diagn. Lab. Immunol.* 3: 444-50); pneumococcal surface protein A (PspA) (McDaniel *et al.*  
 (1997) *Gene Ther.* 4: 375-377); *F. tularensis* outer membrane protein FopA (Fulop *et al.* (1996)  
 10 *FEMS Immunol. Med. Microbiol.* 13: 245-247); the major outer membrane protein within strains  
 of the genus *Actinobacillus* (Hartmann *et al.* (1996) *Zentralbl. Bakteriolog.* 284: 255-262); p60 or  
 listeriolysin (Hly) antigen of *Listeria monocytogenes* (Hess *et al.* (1996) *Proc. Nat'l. Acad. Sci.*  
*USA* 93: 1458-1463); flagellar (G) antigens observed on *Salmonella enteritidis* and *S. pullorum*  
 (Holt and Chaubal (1997) *J. Clin. Microbiol.* 35: 1016-1020); *Bacillus anthracis* protective  
 antigen (PA) (Ivins *et al.* (1995) *Vaccine* 13: 1779-1784); *Echinococcus granulosus* antigen 5  
 (Jones *et al.* (1996) *Parasitology* 113: 213-222); the *rol* genes of *Shigella dysenteriae* 1 and  
*Escherichia coli* K-12 (Klee *et al.* (1997) *J. Bacteriol.* 179: 2421-2425); cell surface proteins Rib  
 and alpha of group B streptococcus (Larsson *et al.* (1996) *Infect. Immun.* 64: 3518-3523); the 37  
 kDa secreted polypeptide encoded on the 70 kb virulence plasmid of pathogenic *Yersinia* spp.  
 (Leary *et al.* (1995) *Contrib. Microbiol. Immunol.* 13: 216-217 and Roggenkamp *et al.* (1997)  
 20 *Infect. Immun.* 65: 446-51); the OspA (outer surface protein A) of the Lyme disease spirochete  
*Borrelia burgdorferi* (Li *et al.* (1997) *Proc. Nat'l. Acad. Sci. USA* 94: 3584-3589, Padilla *et al.*  
 (1996) *J. Infect. Dis.* 174: 739-746, and Wallich *et al.* (1996) *Infection* 24: 396-397); the  
*Brucella melitensis* group 3 antigen gene encoding Omp28 (Lindler *et al.* (1996) *Infect. Immun.*  
 25 64: 2490-2499); the PAc antigen of *Streptococcus mutans* (Murakami *et al.* (1997) *Infect.*  
*Immun.* 65: 794-797); pneumolysin, Pneumococcal neuraminidases, autolysin, hyaluronidase,  
 and the 37 kDa pneumococcal surface adhesin A (Paton *et al.* (1997) *Microb. Drug Resist.* 3: 1-  
 10); 29-32, 41-45, 63-71 x 10(3) MW antigens of *Salmonella typhi* (Perez *et al.* (1996)  
*Immunology* 89: 262-267); K-antigen as a marker of *Klebsiella pneumoniae* (Priamukhina and  
 30 Morozova (1996) *Klin. Lab. Diagn.* 47-9); nocardial antigens of molecular mass approximately  
 60, 40, 20 and 15-10 kDa (Prokesova *et al.* (1996) *Int. J. Immunopharmacol.* 18: 661-668);

*Staphylococcus aureus* antigen ORF-2 (Rieneck *et al.* (1997) *Biochim Biophys Acta* 1350: 128-132); GlpQ antigen of *Borrelia hermsii* (Schwan *et al.* (1996) *J. Clin. Microbiol.* 34: 2483-2492); cholera protective antigen (CPA) (Sciortino (1996) *J. Diarrhoeal Dis. Res.* 14: 16-26); a 190-kDa protein antigen of *Streptococcus mutans* (Senpuku *et al.* (1996) *Oral Microbiol. Immunol.* 11: 121-128); Anthrax toxin protective antigen (PA) (Sharma *et al.* (1996) *Protein Expr. Purif.* 7: 33-38); *Clostridium perfringens* antigens and toxoid (Strom *et al.* (1995) *Br. J. Rheumatol.* 34: 1095-1096); the SEF14 fimbrial antigen of *Salmonella enteritidis* (Thorns *et al.* (1996) *Microb. Pathog.* 20: 235-246); the *Yersinia pestis* capsular antigen (F1 antigen) (Titball *et al.* (1997) *Infect. Immun.* 65: 1926-1930); a 35-kilodalton protein of *Mycobacterium leprae* (Triccas *et al.* (1996) *Infect. Immun.* 64: 5171-5177); the major outer membrane protein, CD, extracted from *Moraxella* (Branhamella) catarrhalis (Yang *et al.* (1997) *FEMS Immunol. Med. Microbiol.* 17: 187-199); pH6 antigen (PsaA protein) of *Yersinia pestis* (Zav'yalov *et al.* (1996) *FEMS Immunol. Med. Microbiol.* 14: 53-57); a major surface glycoprotein, gp63, of *Leishmania major* (Xu and Liew (1994) *Vaccine* 12: 1534-1536; Xu and Liew (1995) *Immunology* 84: 173-176); mycobacterial heat shock protein 65, mycobacterial antigen (*Mycobacterium leprae* hsp65) (Lowrie *et al.* (1994) *Vaccine* 12: 1537-1540; Ragno *et al.* (1997) *Arthritis Rheum.* 40: 277-283; Silva (1995) *Braz. J. Med. Biol. Res.* 28: 843-851); *Mycobacterium tuberculosis* antigen 85 (Ag85) (Huygen *et al.* (1996) *Nat. Med.* 2: 893-898); the 45/47 kDa antigen complex (APA) of *Mycobacterium tuberculosis*, *M. bovis* and BCG (Horn *et al.* (1996) *J. Immunol. Methods* 197: 151-159); the mycobacterial antigen, 65-kDa heat shock protein, hsp65 (Tascon *et al.* (1996) *Nat. Med.* 2: 888-892); the mycobacterial antigens MPB64, MPB70, MPB57 and alpha antigen (Yamada *et al.* (1995) *Kekkaku* 70: 639-644); the *M. tuberculosis* 38 kDa protein (Vordermeier *et al.* (1995) *Vaccine* 13: 1576-1582); the MPT63, MPT64 and MPT-59 antigens from *Mycobacterium tuberculosis* (Manca *et al.* (1997) *Infect. Immun.* 65: 16-23; Oettinger *et al.* (1997) *Scand. J. Immunol.* 45: 499-503; Wilcke *et al.* (1996) *Tuber. Lung Dis.* 77: 250-256); the 35-kilodalton protein of *Mycobacterium leprae* (Triccas *et al.* (1996) *Infect. Immun.* 64: 5171-5177); the ESAT-6 antigen of virulent mycobacteria (Brandt *et al.* (1996) *J. Immunol.* 157: 3527-3533; Pollock and Andersen (1997) *J. Infect. Dis.* 175: 1251-1254); *Mycobacterium tuberculosis* 16-kDa antigen (Hsp16.3) (Chang *et al.* (1996) *J. Biol. Chem.* 271: 7218-7223); and the 18-kilodalton protein of *Mycobacterium leprae* (Baumgart *et al.* (1996) *Infect. Immun.* 64: 2274-2281); protective antigen (PA) of *B. anthracis*; V antigen from *Yersinia pestis*, Y.

*enterocolitica*, and *Y. pseudotuberculosis*; antigens against bacterium *Vibrio cholerae*, cholera toxin B subunit, and heat-labile enterotoxins (LT) from enterotoxigenic *E. coli* strains.

### Viral pathogens

Polypeptides or proteins corresponding to or associated with various viral pathogens, including, but not limited to, e.g., hanta virus (e.g., hanta virus glycoproteins), flaviviruses, such as, e.g., Dengue viruses (e.g., envelope proteins), Japanese, St. Louis and Murray Valley encephalitis viruses, tick-borne encephalitis viruses can be modified according to the invention.

Viral antigens that can be modified according to the invention include, but are not limited to, influenza A virus N2 neuraminidase (Kilbourne *et al.* (1995) *Vaccine* 13: 1799-1803); Dengue virus envelope (E) and premembrane (prM) antigens (Feighny *et al.* (1994) *Am. J. Trop. Med. Hyg.* 50: 322-328; Putnak *et al.* (1996) *Am. J. Trop. Med. Hyg.* 55: 504-10); HIV antigens Gag, Pol, Vif and Nef (Vogt *et al.* (1995) *Vaccine* 13: 202-208); HIV antigens gp120 and gp160 (Achour *et al.* (1995) *Cell. Mol. Biol.* 41: 395-400; Hone *et al.* (1994) *Dev. Biol. Stand.* 82: 159-162); gp41 epitope of human immunodeficiency virus (Eckhart *et al.* (1996) *J. Gen. Virol.* 77: 2001-2008); rotavirus antigen VP4 (Mattion *et al.* (1995) *J. Virol.* 69: 5132-5137); the rotavirus protein VP7 or VP7sc (Emslie *et al.* (1995) *J. Virol.* 69: 1747-1754; Xu *et al.* (1995) *J. Gen. Virol.* 76: 1971-1980); herpes simplex virus (HSV) glycoproteins gB, gC, gD, gE, gG, gH, and gI (Fleck *et al.* (1994) *Med. Microbiol. Immunol.* (Berl) 183: 87-94 [Mattion, 1995]; Ghiasi *et al.* (1995) *Invest. Ophthalmol. Vis. Sci.* 36: 1352-1360; McLean *et al.* (1994) *J. Infect. Dis.* 170: 1100-1109); immediate-early protein ICP47 of herpes simplex virus-type 1 (HSV-1) (Banks *et al.* (1994) *Virology* 200: 236-245); immediate-early (IE) proteins ICP27, ICP0, and ICP4 of herpes simplex virus (Manickan *et al.* (1995) *J. Virol.* 69: 4711-4716); influenza virus nucleoprotein and hemagglutinin (Deck *et al.* (1997) *Vaccine* 15: 71-78; Fu *et al.* (1997) *J. Virol.* 71: 2715-2721); B19 parvovirus capsid proteins VP1 (Kawase *et al.* (1995) *Virology* 211: 359-366) or VP2 (Brown *et al.* (1994) *Virology* 198: 477-488); Hepatitis B virus core and e antigen and capsid protein (Schodel *et al.* (1996) *Intervirology* 39: 104-106); hepatitis B surface antigen (Shiau and Murray (1997) *J. Med. Virol.* 51: 159-166); hepatitis B surface antigen fused to the core antigen of the virus (Id.); Hepatitis B virus core-preS2 particles (Nemeckova *et al.* (1996) *Acta Virol.* 40: 273-279); HBV preS2-S protein (Kutinova *et al.* (1996) *Vaccine* 14: 1045-1052); VZV glycoprotein I (Kutinova *et al.* (1996) *Vaccine* 14: 1045-1052); rabies virus glycoproteins

(Xiang *et al.* (1994) *Virology* 199: 132-140; Xuan *et al.* (1995) *Virus Res.* 36: 151-161) or ribonucleocapsid (Hooper *et al.* (1994) *Proc. Nat'l. Acad. Sci. USA* 91: 10908-10912); human cytomegalovirus (HCMV) glycoprotein B (UL55) (Britt *et al.* (1995) *J. Infect. Dis.* 171: 18-25); the hepatitis C virus (HCV) nucleocapsid protein in a secreted or a nonsecreted form, or as a fusion protein with the middle (pre-S2 and S) or major (S) surface antigens of hepatitis B virus (HBV) (Inchauspe *et al.* (1997) *DNA Cell Biol.* 16: 185-195; Major *et al.* (1995) *J. Virol.* 69: 5798-5805); the hepatitis C virus antigens: the core protein (pC); E1 (pE1) and E2 (pE2) alone or as fusion proteins (Saito *et al.* (1997) *Gastroenterology* 112: 1321-1330); the gene encoding respiratory syncytial virus fusion protein (PFP-2) (Falsey and Walsh (1996) *Vaccine* 14: 1214-1218; Piedra *et al.* (1996) *Pediatr. Infect. Dis. J.* 15: 23-31); the VP6 and VP7 genes of rotaviruses (Choi *et al.* (1997) *Virology* 232: 129-138; Jin *et al.* (1996) *Arch. Virol.* 141: 2057-2076); the E1, E2, E3, E4, E5, E6 and E7 proteins of human papillomavirus (Brown *et al.* (1994) *Virology* 201: 46-54; Dillner *et al.* (1995) *Cancer Detect. Prev.* 19: 381-393; Krul *et al.* (1996) *Cancer Immunol. Immunother.* 43: 44-48; Nakagawa *et al.* (1997) *J. Infect. Dis.* 175: 927-931); a human T-lymphotropic virus type I gag protein (Porter *et al.* (1995) *J. Med. Virol.* 45: 469-474); Epstein-Barr virus (EBV) gp340 (Mackett *et al.* (1996) *J. Med. Virol.* 50: 263-271); the Epstein-Barr virus (EBV) latent membrane protein LMP2 (Lee *et al.* (1996) *Eur. J. Immunol.* 26: 1875-1883); Epstein-Barr virus nuclear antigens 1 and 2 (Chen and Cooper (1996) *J. Virol.* 70: 4849-4853; Khanna *et al.* (1995) *Virology* 214: 633-637); the measles virus nucleoprotein (N) (Fooks *et al.* (1995) *Virology* 210: 456-465); and cytomegalovirus glycoprotein gB (Marshall *et al.* (1994) *J. Med. Virol.* 43: 77-83) or glycoprotein gH (Rasmussen *et al.* (1994) *J. Infect. Dis.* 170: 673-677).

### Parasites

Antigens from parasites can also be modified according to the invention. These include, but are not limited to, the schistosome gut-associated antigens CAA (circulating anodic antigen) and CCA (circulating cathodic antigen) in *Schistosoma mansoni*, *S. haematobium* or *S. japonicum* (Deelder *et al.* (1996) *Parasitology* 112: 21-35); a multiple antigen peptide (MAP) composed of two distinct protective antigens derived from the parasite *Schistosoma mansoni* (Ferru *et al.* (1997) *Parasite Immunol.* 19: 1-11); *Leishmania* parasite surface molecules (Lezama-Davila (1997) *Arch. Med. Res.* 28: 47-53); third-stage larval (L3) antigens of *L. loa* (Akue *et al.* (1997) *J. Infect. Dis.* 175: 158-63); the genes, Tams1-1 and Tams1-2, encoding the

30-and 32-kDa major merozoite surface antigens of *Theileria annulata* (Ta) (d'Oliveira *et al.* (1996) *Gene* 172: 33-39); *Plasmodium falciparum* merozoite surface antigen 1 or 2 (al-Yaman *et al.* (1995) *Trans. R. Soc. Trop. Med. Hyg.* 89: 555-559; Beck *et al.* (1997) *J. Infect. Dis.* 175: 921-926; Rzepczyk *et al.* (1997) *Infect. Immun.* 65: 1098-1100); circumsporozoite (CS) protein-based B-epitopes from *Plasmodium berghei*, (PPPPNPND)<sub>2</sub> and *Plasmodium yoelii*, (QGPGAP)<sub>3</sub>QG, along with a *P. berghei* T-helper epitope KQIRDSITEEWS (Reed *et al.* (1997) *Vaccine* 15: 482-488); NYVAC-Pf7 encoded *Plasmodium falciparum* antigens derived from the sporozoite (circumsporozoite protein and sporozoite surface protein 2), liver (liver stage antigen 1), blood (merozoite surface protein 1, serine repeat antigen, and apical membrane antigen 1), and sexual (25-kDa sexual-stage antigen) stages of the parasite life cycle were inserted into a single NYVAC genome to generate NYVAC-Pf7 (Tine *et al.* (1996) *Infect. Immun.* 64: 3833-3844); *Plasmodium falciparum* antigen Pfs230 (Williamson *et al.* (1996) *Mol. Biochem. Parasitol.* 78: 161-169); *Plasmodium falciparum* apical membrane antigen (AMA-1) (Lal *et al.* (1996) *Infect. Immun.* 64: 1054-1059); *Plasmodium falciparum* proteins Pfs28 and Pfs25 (Duffy and Kaslow (1997) *Infect. Immun.* 65: 1109-1113); *Plasmodium falciparum* merozoite surface protein, MSP1 (Hui *et al.* (1996) *Infect. Immun.* 64: 1502-1509); the malaria antigen Pf332 (Ahlborg *et al.* (1996) *Immunology* 88: 630-635); *Plasmodium falciparum* erythrocyte membrane protein 1 (Baruch *et al.* (1995) *Proc. Nat'l. Acad. Sci. USA* 93: 3497-3502; Baruch *et al.* (1995) *Cell* 82: 77-87); *Plasmodium falciparum* merozoite surface antigen, PfMSP-1 (Egan *et al.* (1996) *J. Infect. Dis.* 173: 765-769); *Plasmodium falciparum* antigens SERA, EBA-175, RAP1 and RAP2 (Riley (1997) *J. Pharm. Pharmacol.* 49: 21-27); *Schistosoma japonicum* paramyosin (Sj97) or fragments thereof (Yang *et al.* (1995) *Biochem. Biophys. Res. Commun.* 212: 1029-1039); and Hsp70 in parasites (Maresca and Kobayashi (1994) *Experientia* 50: 1067-1074).

### Allergen antigens

Allergen antigens that can be modified according to the invention, include, but are not limited to those of animals, including the mite (*e.g.*, *Dermatophagoides pteronyssinus*, *Dermatophagoides farinae*, *Blomia tropicalis*), such as the allergens der p1 (Scobie *et al.* (1994) *Biochem. Soc. Trans.* 22: 448S; Yssel *et al.* (1992) *J. Immunol.* 148: 738-745), der p2 (Chua *et al.* (1996) *Clin. Exp. Allergy* 26: 829-837), der p3 (Smith and Thomas (1996) *Clin. Exp. Allergy* 26: 571-579), der p5, der p V (Lin *et al.* (1994) *J. Allergy Clin. Immunol.* 94: 989-996), der p6 (Bennett and Thomas (1996) *Clin. Exp. Allergy* 26: 1150-1154), der p 7 (Shen *et al.* (1995) *Clin.*

*Exp. Allergy* 25: 416-422), der f2 (Yuuki *et al.* (1997) *Int. Arch. Allergy Immunol.* 112: 44-48), der f3 (Nishiyama *et al.* (1995) *FEBS Lett.* 377: 62-66), der f7 (Shen *et al.* (1995) *Clin. Exp. Allergy* 25: 1000-1006); Mag 3 (Fujikawa *et al.* (1996) *Mol. Immunol.* 33: 311-319). Also of interest as antigens are the house dust mite allergens Tyr p2 (Eriksson *et al.* (1998) *Eur. J. Biochem.* 251: 443-447), Lep d1 (Schmidt *et al.* (1995) *FEBS Lett.* 370: 11-14), and glutathione S-transferase (O'Neill *et al.* (1995) *Immunol Lett.* 48: 103-107); the 25,589 Da, 219 amino acid polypeptide with homology with glutathione S-transferases (O'Neill *et al.* (1994) *Biochim. Biophys. Acta.* 1219: 521-528); Blo t 5 (Arruda *et al.* (1995) *Int. Arch. Allergy Immunol.* 107: 456-457); bee venom phospholipase A2 (Carballido *et al.* (1994) *J. Allergy Clin. Immunol.* 93: 758-767; Jutel *et al.* (1995) *J. Immunol.* 154: 4187-4194); bovine dermal/dander antigens BDA 11 (Rautiainen *et al.* (1995) *J. Invest. Dermatol.* 105: 660-663) and BDA20 (Mantylarvi *et al.* (1996) *J. Allergy Clin. Immunol.* 97: 1297-1303); the major horse allergen Equ c1 (Gregoire *et al.* (1996) *J. Biol. Chem.* 271: 32951-32959); Jumper ant *M. pilosula* allergen Myr p I and its homologous allergenic polypeptides Myr p2 (Donovan *et al.* (1996) *Biochem. Mol. Biol. Int.* 39: 877-885); 1-13, 14, 16 kD allergens of the mite *Blomia tropicalis* (Caraballo *et al.* (1996) *J. Allergy Clin. Immunol.* 98: 573-579); the cockroach allergens Bla g Bd90K (Helm *et al.* (1996) *J. Allergy Clin. Immunol.* 98: 172-80) and Bla g 2 (Arruda *et al.* (1995) *J. Biol. Chem.* 270: 19563-19568); the cockroach Cr-PI allergens (Wu *et al.* (1996) *J. Biol. Chem.* 271: 17937-17943); fire ant venom allergen, Sol i 2 (Schmidt *et al.* (1996) *J. Allergy Clin. Immunol.* 98: 82-88); the insect *Chironomus thummi* major allergen Chi t 1-9 (Kipp *et al.* (1996) *Int. Arch. Allergy Immunol.* 110: 348-353); dog allergen Can f 1 or cat allergen Fel d 1 (Ingram *et al.* (1995) *J. Allergy Clin. Immunol.* 96: 449-456); albumin, derived, for example, from horse, dog or cat (Goubran Botros *et al.* (1996) *Immunology* 88: 340-347); deer allergens with the molecular mass of 22 kD, 25 kD or 60 kD (Spitzauer *et al.* (1997) *Clin. Exp. Allergy* 27: 196-200); and the 20 kd major allergen of cow (Ylonen *et al.* (1994) *J. Allergy Clin. Immunol.* 93: 851-858).

Pollen and grass allergens can also be modified according to the invention. Such allergens include, for example, Hor v9 (Astwood and Hill (1996) *Gene* 182: 53-62, Lig v1 (Batanero *et al.* (1996) *Clin. Exp. Allergy* 26: 1401-1410); Lol p 1 (Muller *et al.* (1996) *Int. Arch. Allergy Immunol.* 109: 352-355), Lol p II (Tamborini *et al.* (1995) *Mol. Immunol.* 32: 505-513), Lol pVA, Lol pVB (Ong *et al.* (1995) *Mol. Immunol.* 32: 295-302), Lol p 9 (Blaher *et al.* (1996) *J. Allergy Clin. Immunol.* 98: 124-132); Par J I (Costa *et al.* (1994) *FEBS Lett.* 341: 182-186;

Sallusto *et al.* (1996) *J. Allergy Clin. Immunol.* 97: 627-637), Par j 2.0101 (Duro *et al.* (1996) *FEBS Lett.* 399: 295-298); Bet v1 (Faber *et al.* (1996) *J. Biol. Chem.* 271: 19243-19250), Bet v2 (Rihs *et al.* (1994) *Int. Arch. Allergy Immunol.* 105: 190-194); Dac g3 (Guerin-Marchand *et al.* (1996) *Mol. Immunol.* 33: 797-806); Phl p 1 (Petersen *et al.* (1995) *J. Allergy Clin. Immunol.* 95: 987-994), Phl p 5 (Muller *et al.* (1996) *Int. Arch. Allergy Immunol.* 109: 352-355), Phl p 6 (Petersen *et al.* (1995) *Int. Arch. Allergy Immunol.* 108: 55-59); Cry j I (Sone *et al.* (1994) *Biochem. Biophys. Res. Commun.* 199: 619-625), Cry j II (Namba *et al.* (1994) *FEBS Lett.* 353: 124-128); Cor a 1 (Schenk *et al.* (1994) *Eur. J. Biochem.* 224: 717-722); cyn d1 (Smith *et al.* (1996) *J. Allergy Clin. Immunol.* 98: 331-343), cyn d7 (Suphioglu *et al.* (1997) *FEBS Lett.* 402: 167-172); Pha a 1 and isoforms of Pha a 5 (Suphioglu and Singh (1995) *Clin. Exp. Allergy* 25: 853-865); Cha o 1 (Suzuki *et al.* (1996) *Mol. Immunol.* 33: 451-460); profilin derived, i.e., from timothy grass or birch pollen (Valenta *et al.* (1994) *Biochem. Biophys. Res. Commun.* 199: 106-118); P0149 (Wu *et al.* (1996) *Plant Mol. Biol.* 32: 1037-1042); Ory s1 (Xu *et al.* (1995) *Gene* 164: 255-259); and Amb a V and Amb t 5 (Kim *et al.* (1996) *Mol. Immunol.* 33: 873-880; Zhu *et al.* (1995) *J. Immunol.* 155: 5064-5073).

Food allergens that can be modified according to the invention include, for example, profilin (Rihs *et al.* (1994) *Int. Arch. Allergy Immunol.* 105: 190-194); rice allergenic cDNAs belonging to the alpha-amylase/trypsin inhibitor gene family (Alvarez *et al.* (1995) *Biochim Biophys Acta* 1251: 201-204); the main olive allergen, Ole e I (Lombardero *et al.* (1994) *Clin Exp Allergy* 24: 765-770); Sin a 1, the major allergen from mustard (Gonzalez De La Pena *et al.* (1996) *Eur J Biochem.* 237: 827-832); parvalbumin, the major allergen of salmon (Lindstrom *et al.* (1996) *Scand. J. Immunol.* 44: 335-344); apple allergens, such as the major allergen Mal d 1 (Vanek-Krebitz *et al.* (1995) *Biochem. Biophys. Res. Commun.* 214: 538-551); and peanut allergens, such as Ara h I (Burks *et al.* (1995) *J. Clin. Invest.* 96: 1715-1721).

Fungal allergens that can be modified according to the invention include, but are not limited to, the allergen, Cla h III, of *Cladosporium herbarum* (Zhang *et al.* (1995) *J. Immunol.* 154: 710-717); the allergen Psi c 2, a fungal cyclophilin, from the basidiomycete *Psilocybe cubensis* (Horner *et al.* (1995) *Int. Arch. Allergy Immunol.* 107: 298-300); hsp 70 cloned from a cDNA library of *Cladosporium herbarum* (Zhang *et al.* (1996) *Clin Exp Allergy* 26: 88-95); the 68 kD allergen of *Penicillium notatum* (Shen *et al.* (1995) *Clin. Exp. Allergy* 26: 350-356); aldehyde dehydrogenase (ALDH) (Achatz *et al.* (1995) *Mol Immunol.* 32: 213-227);

enolase (Achatz *et al.* (1995) *Mol. Immunol.* 32: 213-227); YCP4 (Id.); acidic ribosomal protein P2 (Id.).

Other allergens that can be modified include latex allergens, such as a major allergen (Hev b 5) from natural rubber latex (Akasawa *et al.* (1996) *J. Biol. Chem.* 271: 25389-25393; Slater *et al.* (1996) *J. Biol. Chem.* 271: 25394-25399).

Antigens associated with autoimmune diseases and inflammatory conditions

Autoantigens that can be modified according to the invention include, but are not limited to, myelin basic protein (Stinissen *et al.* (1996) *J. Neurosci. Res.* 45: 500-511) or a fusion protein of myelin basic protein and proteolipid protein (Elliott *et al.* (1996) *J. Clin. Invest.* 98: 1602-1612), proteolipid protein (PLP) (Rosener *et al.* (1997) *J. Neuroimmunol.* 75: 28-34), 2',3'-cyclic nucleotide 3'-phosphodiesterase (CNPase) (Rosener *et al.* (1997) *J. Neuroimmunol.* 75: 28-34), the Epstein Barr virus nuclear antigen-1 (EBNA-1) (Vaughan *et al.* (1996) *J. Neuroimmunol.* 69: 95-102), HSP70 (Salveti *et al.* (1996) *J. Neuroimmunol.* 65: 143-53; Feldmann *et al.* (1996) *Cell* 85: 307).

Antigens that can be modified according to the invention and used to treat scleroderma, systemic sclerosis, and systemic lupus erythematosus include, for example, (-2-GPI, 50 kDa glycoprotein (Blank *et al.* (1994) *J. Autoimmun.* 7: 441-455), Ku (p70/p80) autoantigen, or its 80-kd subunit protein (Hong *et al.* (1994) *Invest. Ophthalmol. Vis. Sci.* 35: 4023-4030; Wang *et al.* (1994) *J. Cell Sci.* 107: 3223-3233), the nuclear autoantigens La (SS-B) and Ro (SS-A) (Huang *et al.* (1997) *J. Clin. Immunol.* 17: 212-219; Igarashi *et al.* (1995) *Autoimmunity* 22: 33-42; Keech *et al.* (1996) *Clin. Exp. Immunol.* 104: 255-263; Manoussakis *et al.* (1995) *J. Autoimmun.* 8: 959-969; Topfer *et al.* (1995) *Proc. Nat'l. Acad. Sci. USA* 92: 875-879), proteasome (-type subunit C9 (Feist *et al.* (1996) *J. Exp. Med.* 184: 1313-1318), Scleroderma antigens Rpp 30, Rpp 38 or Scl-70 (Eder *et al.* (1997) *Proc. Nat'l. Acad. Sci. USA* 94: 1101-1106; Hietarinta *et al.* (1994) *Br. J. Rheumatol.* 33: 323-326), the centrosome autoantigen PCM-1 (Bao *et al.* (1995) *Autoimmunity* 22: 219-228), polymyositis-scleroderma autoantigen (PM-Scl) (Kho *et al.* (1997) *J. Biol. Chem.* 272: 13426-13431), scleroderma (and other systemic autoimmune disease) autoantigen CENP-A (Muro *et al.* (1996) *Clin. Immunol. Immunopathol.* 78: 86-89), U5, a small nuclear ribonucleoprotein (snRNP) (Okano *et al.* (1996) *Clin. Immunol. Immunopathol.* 81: 41-47), the 100-kd protein of PM-Scl autoantigen (Ge *et al.* (1996) *Arthritis Rheum.* 39: 1588-1595), the nucleolar U3- and Th(7-2) ribonucleoproteins

(Verheijen *et al.* (1994) *J. Immunol. Methods* 169: 173-182), the ribosomal protein L7 (Neu *et al.* (1995) *Clin. Exp. Immunol.* 100: 198-204), hPop1 (Lygerou *et al.* (1996) *EMBO J.* 15: 5936-5948), and a 36-kd protein from nuclear matrix antigen (Deng *et al.* (1996) *Arthritis Rheum.* 39: 1300-1307).

5           Antigens useful in treatment of hepatic autoimmune disorders can also be modified; these include the cytochromes P450 and UDP-glucuronosyl-transferases (Obermayer-Straub and Manns (1996) *Baillieres Clin. Gastroenterol.* 10: 501-532), the cytochromes P450 2C9 and P450 1A2 (Bourdi *et al.* (1996) *Chem. Res. Toxicol.* 9: 1159-1166; Clemente *et al.* (1997) *J. Clin. Endocrinol. Metab.* 82: 1353-1361), LC-1 antigen (Klein *et al.* (1996) *J. Pediatr. Gastroenterol. Nutr.* 23: 461-465), and a 230-kDa Golgi-associated protein (Funaki *et al.* (1996) *Cell Struct. Funct.* 21: 63-72).

10           Antigens useful for treatment of autoimmune disorders of the skin that can be modified according to the invention include, but are not limited to, the 450 kD human epidermal autoantigen (Fujiwara *et al.* (1996) *J. Invest. Dermatol.* 106: 1125-1130), the 230 kD and 180 kD bullous pemphigoid antigens (Hashimoto (1995) *Keio J. Med.* 44: 115-123; Murakami *et al.* (1996) *J. Dermatol. Sci.* 13: 112-117), pemphigus foliaceus antigen (desmoglein 1), pemphigus vulgaris antigen (desmoglein 3), BPAg2, BPAg1, and type VII collagen (Batteux *et al.* (1997) *J. Clin. Immunol.* 17: 228-233; Hashimoto *et al.* (1996) *J. Dermatol. Sci.* 12: 10-17), a 168-kDa mucosal antigen in a subset of patients with cicatricial pemphigoid (Ghohestani *et al.* (1996) *J. Invest. Dermatol.* 107: 136-139), and a 218-kd nuclear protein (218-kd Mi-2) (Seelig *et al.* (1995) *Arthritis Rheum.* 38: 1389-1399).

15           Antigens for treating insulin dependent diabetes mellitus can also be modified; these, include, but are not limited to, insulin, proinsulin, GAD65 and GAD67, heat-shock protein 65 (hsp65), and islet-cell antigen 69 (ICA69) (French *et al.* (1997) *Diabetes* 46: 34-39; Roep (1996) *Diabetes* 45: 1147-1156; Schloot *et al.* (1997) *Diabetologia* 40: 332-338), viral proteins homologous to GAD65 (Jones and Crosby (1996) *Diabetologia* 39: 1318-1324), islet cell antigen-related protein-tyrosine phosphatase (PTP) (Cui *et al.* (1996) *J. Biol. Chem.* 271: 24817-24823), GM2-1 ganglioside (Cavallo *et al.* (1996) *J. Endocrinol.* 150: 113-120; Dotta *et al.* (1996) *Diabetes* 45: 1193-1196), glutamic acid decarboxylase (GAD) (Nepom (1995) *Curr. Opin. Immunol.* 7: 825-830; Panina-Bordignon *et al.* (1995) *J. Exp. Med.* 181: 1923-1927), an islet cell antigen (ICA69) (Karges *et al.* (1997) *Biochim. Biophys. Acta* 1360: 97-101; Roep *et al.*

(1996) *Eur. J. Immunol.* 26: 1285-1289), Tep69, the single T cell epitope recognized by T cells from diabetes patients (Karges *et al.* (1997) *Biochim. Biophys. Acta* 1360: 97-101), ICA 512, an autoantigen of type I diabetes (Solimena *et al.* (1996) *EMBO J.* 15: 2102-2114), an islet-cell protein tyrosine phosphatase and the 37-kDa autoantigen derived from it in type 1 diabetes (including IA-2, IA-2) (La Gasse *et al.* (1997) *Mol. Med.* 3: 163-173), the 64 kDa protein from In-111 cells or human thyroid follicular cells that is immunoprecipitated with sera from patients with islet cell surface antibodies (ICSA) (Igawa *et al.* (1996) *Endocr. J.* 43: 299-306), phogrin, a homologue of the human transmembrane protein tyrosine phosphatase, an autoantigen of type 1 diabetes (Kawasaki *et al.* (1996) *Biochem. Biophys. Res. Commun.* 227: 440-447), the 40 kDa and 37 kDa tryptic fragments and their precursors IA-2 and IA-2 in IDDM (Lampasona *et al.* (1996) *J. Immunol.* 157: 2707-2711; Notkins *et al.* (1996) *J. Autoimmun.* 9: 677-682), insulin or a cholera toxoid-insulin polypeptide (Bergerot *et al.* (1997) *Proc. Nat'l. Acad. Sci. USA* 94: 4610-4614), carboxypeptidase H, the human homologue of gp330, which is a renal epithelial glycoprotein involved in inducing Heymann nephritis in rats, and the 38-kD islet mitochondrial autoantigen (Arden *et al.* (1996) *J. Clin. Invest.* 97: 551-561).

Useful antigens for rheumatoid arthritis treatment that can be modified according to the invention include, but are not limited to, the 45 kDa DEK nuclear antigen, in particular onset juvenile rheumatoid arthritis and iridocyclitis (Murray *et al.* (1997) *J. Rheumatol.* 24: 560-567), human cartilage glycoprotein-39, an autoantigen in rheumatoid arthritis (Verheijden *et al.* (1997) *Arthritis Rheum.* 40: 1115-1125), a 68k autoantigen in rheumatoid arthritis (Blass *et al.* (1997) *Ann. Rheum. Dis.* 56: 317-322), collagen (Rosloniec *et al.* (1995) *J. Immunol.* 155: 4504-4511), collagen type II (Cook *et al.* (1996) *Arthritis Rheum.* 39: 1720-1727; Trentham (1996) *Ann. N. Y. Acad. Sci.* 778: 306-314), cartilage link protein (Guerassimov *et al.* (1997) *J. Rheumatol.* 24: 959-964), ezrin, radixin and moesin, which are auto-immune antigens in rheumatoid arthritis (Wagatsuma *et al.* (1996) *Mol. Immunol.* 33: 1171-1176), and mycobacterial heat shock protein 65 (Ragno *et al.* (1997) *Arthritis Rheum.* 40: 277-283).

Antigens useful for treatment are autoimmune thyroid disorders that can be modified include, for example, thyroid peroxidase and the thyroid stimulating hormone receptor (Tandon and Weetman (1994) *J. R. Coll. Physicians Lond.* 28: 10-18), thyroid peroxidase from human Graves' thyroid tissue (Gardas *et al.* (1997) *Biochem. Biophys. Res. Commun.* 234: 366-370; Zimmer *et al.* (1997) *Histochem. Cell. Biol.* 107: 115-120), a 64-kDa antigen associated

with thyroid-associated ophthalmopathy (Zhang *et al.* (1996) *Clin. Immunol. Immunopathol.* 80: 236-244), the human TSH receptor (Nicholson *et al.* (1996) *J. Mol. Endocrinol.* 16: 159-170), and the 64 kDa protein from In-111 cells or human thyroid follicular cells that is immunoprecipitated with sera from patients with islet cell surface antibodies (ICSA) (Igawa *et al.* (1996) *Endocr. J.* 43: 299-306).

Other associated antigens that can be modified include, but are not limited to, Sjogren's syndrome (-fodrin; Haneji *et al.* (1997) *Science* 276: 604-607), myasthenia gravis (the human M2 acetylcholine receptor or fragments thereof, specifically the second extracellular loop of the human M2 acetylcholine receptor; Fu *et al.* (1996) *Clin. Immunol. Immunopathol.* 78: 203-207), vitiligo (tyrosinase; Fishman *et al.* (1997) *Cancer* 79: 1461-1464), a 450 kD human epidermal autoantigen recognized by serum from individual with blistering skin disease, and ulcerative colitis (chromosomal proteins HMG1 and HMG2; Sobajima *et al.* (1997) *Clin. Exp. Immunol.* 107: 135-140).

#### Sperm Antigens

Sperm antigens which can be used in the genetic vaccines include, for example, lactate dehydrogenase (LDH-C4), galactosyltransferase (GT), SP-10, rabbit sperm autoantigen (RSA), guinea pig (g)PH-20, cleavage signal protein (CS-1), HSA-63, human (h)PH-20, and AgX-1 (Zhu and Naz (1994) *Arch. Androl.* 33: 141-144), the synthetic sperm peptide, P10G (O'Rand *et al.* (1993) *J. Reprod. Immunol.* 25: 89-102), the 135kD, 95kD, 65kD, 47kD, 41kD and 23kD proteins of sperm, and the FA-1 antigen (Naz *et al.* (1995) *Arch. Androl.* 35: 225-231), and the 35 kD fragment of cytokeratin 1 (Lucas *et al.* (1996) *Anticancer Res.* 16: 2493-2496).

Also, examples of antigens are set forth in Punnonen *et al.* (1999) WO 99/41369; Punnonen *et al.* (1999) WO 99/41383; Punnonen *et al.* (1999) WO 99/41368; and Punnonen *et al.* (1999) WO 99/41402), the contents of all of which are incorporated herein by reference in their entirety for all purposes. Other useful antigens have been described in the literature or can be discovered using genomics approaches.

#### Peptide addition

In principle the peptide addition X can be any stretch of amino acid residues ranging from a single amino acid residue to a large protein, e.g., a mature protein. Usually, the peptide addition X comprises 1-500 amino acid residues, such as 2-500, normally 2-50 or 3-50 amino acid residues, such as 3-20 amino acid residues. The length of the peptide addition to be

used for modification of a given polypeptide is dependent of or determined on the basis of a number of factors including the type of polypeptide of interest and the desired effect to be achieved by the modification. Normally, the peptide addition has less than 90% identity to the amino acid sequence of a native full length polypeptide, in particular less than 80% identity, such as less than 70% identity or even lower degree of identity to a full length protein. In one embodiment, the peptide addition may constitute a part of a full length protein (e.g., 1-50 amino acid residues thereof).

The peptide addition may be designed by a site-specific or random approach, e.g. as out-lined in further detail in the Methods section below. This section also comprises a set of guidelines useful for preparing a peptide addition for use in the present invention are described. It will be understood that those guidelines are intended for illustration purposes only and that a person skilled in the art will be aware of alternative useful routes for design of peptide addition. Thus, the method of designing a peptide addition for use herein should not be considered limited to that described in the Materials section.

The number of glycosylation sites should be sufficient to provide the desired effect. Typically, the peptide addition X comprises 1-20, such as 1-10 glycosylation sites. For instance, the peptide addition X comprises 1, 2, 3, 4, 5, 6, 7, 8, 9 or 10 glycosylation sites. It is well known that one frequently occurring consequence of modifying an amino acid sequence of, e.g., a human protein is that new epitopes are created by such modification. In order to shield any new epitopes created by the peptide addition, it is desirable that sufficient glycosylation sites are present to enable shielding of all epitopes introduced into the sequence. This is e.g., achieved when the peptide addition X comprises at least one glycosylation site within a stretch of 30 contiguous amino acid residues, such as at least one glycosylation site within 20 amino acid residues or at least one glycosylation site within 10 amino acid residues, in particular 1-3 glycosylation sites within a stretch of 10 contiguous amino acid residues in the peptide addition X.

Thus, in one embodiment, the peptide addition X comprises at least two glycosylation sites, wherein two of said sites are separated by at most 10 amino acid residues, none of which comprises a glycosylation site. Furthermore, the polypeptide Pp can comprise at least one introduced glycosylation site, in particular 1-5 introduced glycosylation sites.

Analogously, the polypeptide Pp can comprise at least one removed glycosylation site, in particular 1-5 removed glycosylation sites.

The glycosylation site of the peptide addition may be an *in vivo* or *in vitro* glycosylation site. Preferably, the glycosylation site is an *in vivo* glycosylation site, in particular an N-glycosylation site since glycosylation of such site is more easy to control than to an O-glycosylation site. Accordingly, in an embodiment, the peptide addition X comprises at least one N-glycosylation site, typically at least two N-glycosylation sites. For instance, the peptide addition X has the structure  $X_1\text{-N-X}_2\text{-[T/S]/C-Z}$ , wherein  $X_1$  is a peptide comprising at least one amino acid residue or is absent,  $X_2$  is any amino acid residue different from Pro, and Z is absent or a peptide comprising at least one amino acid residue. For instance,  $X_1$  is absent,  $X_2$  is an amino acid residue selected from the group consisting of I, A, G, V and S (all relatively small amino acid residues), and Z comprises at least 1 amino acid residue.

For instance, Z can be a peptide comprising 1-50 amino acid residues and, e.g., 1-10 glycosylation sites.

In another polypeptide of the invention  $X_1$  comprises at least one amino acid residue, e.g., 1-50 amino acid residues,  $X_2$  is an amino acid residue selected from the group consisting of I, A, G, V and S, and Z is absent. For instance,  $X_1$  comprises 1-10 glycosylation sites.

For instance, the peptide addition for use in the present invention can comprise a peptide sequence selected from the group consisting of INA[T/S], GNI[T/S], VNI[T/S], SNI[T/S], ASNI[T/S], NI[T/S], SPINA[T/S], ASPINA[T/S], ANI[T/S]ANI[T/S]ANI, ANI[T/S]GSNI[T/S]GSNI[T/S], FNI[T/S]VNI[T/S]V, YNI[T/S]VNI[T/S]V, AFNI[T/S]VNI[T/S]V, AYNI[T/S]VNI[T/S]V, APND[T/S]VNI[T/S]V, ANI[T/S], ASNS[T/S]NNG[T/S]LNA[T/S], ANH[T/S]NE[T/S]NA[T/S], GSPINA[T/S], ASPINA[T/S]SPINA[T/S], ANN[T/S]NY[T/S]NW[T/S], ATNI[T/S]LNY[T/S]AN[T/S]T, AANS[T/S]GNI[T/S]ING[T/S], AVNW[T/S]SND[T/S]SNS[T/S], GNA[T/S], AVNW[T/S]SND[T/S]SNS[T/S], ANN[T/S]NY[T/S]NS[T/S], ANNTNYTNWT, ANI[T/S]VNI[T/S]V, ND[T/S]VNF[T/S] and NI[T/S]VNI[T/S]V wherein [T/S] is either a T or an S residue, preferably a T residue. Other non-limiting examples include a peptide addition comprising the sequence NSTQNATA, which corresponds to positions 231 to 238 of the human calcium activated channel 2 precursor (to add two N-glycosylation sites), or the sequence

ANLTVRNLTRNVTV, which corresponds to positions 538 to 551 of the human G protein coupled receptor 64 (to add three N-glycosylation sites).

The peptide addition can comprise one or more of these peptide sequences, i.e., at least two of said sequences either directly linked together or separated by one or more amino acid residues, or can contain two or more copies of any of these peptide sequence. It will be understood that the above specific sequences are given for illustrative purposes and thus do not constitute an exclusive list of peptide sequences of use in the present invention.

In a more specific embodiment, the peptide addition X is selected from the group consisting of INA[T/S], GNI[T/S], VNI[T/S], SNI[T/S], ASNI[T/S], NI[T/S], SPINA[T/S], ASPINA[T/S], ANI[T/S]ANI[T/S]ANI, and ANI[T/S]GSNI[T/S]GSNI[T/S], wherein [T/S] is either a T or an S residue, preferably a T residue.

As stated further above the polypeptide Pp can be a native polypeptide that optionally comprises one or more glycosylation sites. In order to further modify the glycosylation of the polypeptide Pp of interest (in terms of the number of oligosaccharide moieties attached to the polypeptide), the polypeptide Pp can be a variant of a native polypeptide that differs from said polypeptide in at least one introduced or at least one removed glycosylation site.

For instance, the polypeptide Pp comprises at least one introduced glycosylation site, in particular 1-5 introduced glycosylation sites, such as 2-5 introduced glycosylation sites.

In order to affect the total glycosylation of the polypeptide of interest the glycosylation site is introduced so that the N residue of said glycosylation site is exposed at the surface of the polypeptide, when folded in its active form. Likewise, a glycosylation site to be removed is selected from those having an N residue exposed at the surface of the polypeptide.

In one embodiment, the peptide addition X has an N residue in position -2 or -1, and the polypeptide Pp or P<sub>x</sub> has a T or an S residue in position +1 or +2, respectively, the residue numbering being made relative to the N-terminal amino acid residue of Pp or P<sub>x</sub>, whereby an N-glycosylation site is formed.

#### Glycosylation

The polypeptide of the invention is glycosylated (i.e., comprises an *in vivo* attached N- or O-linked oligosaccharide moiety or *in vitro* attached oligosaccharide moiety) and furthermore has an altered glycosylation profile as compared to that of the polypeptide Pp. For

instance, the altered glycosylation profile is a consequence of an altered, normally increased, number of attached oligosaccharide moieties and/or an altered type or distribution of attached oligosaccharide moieties.

Furthermore, for polypeptides intended for therapeutic or veterinary uses or to which a human or animal is otherwise exposed, the type of oligosaccharide moiety to be attached should normally be one that does not lead to increased immunogenicity of the polypeptide as compared to that of the polypeptide Pp. The coupling of an oligosaccharide moiety may take place *in vivo* or *in vitro*. In order to achieve *in vivo* glycosylation of a nucleotide sequence encoding the polypeptide should be inserted in a glycosylating, eucaryotic expression host. The expression host cell may be selected from fungal (filamentous fungal or yeast), insect, mammalian cells or transgenic plant cells as disclosed in further detail in the section entitled "Methods of preparing a polypeptide of the invention". Also, the glycosylation may be achieved in the human body when using a nucleotide sequence encoding the polypeptide of the invention in gene therapy.

*In vitro* glycosylation can be achieved by attaching chemically synthesized oligosaccharide structures to the polypeptide using a variety of different chemistries e.g., the chemistries employed for attachment of PEG to proteins, wherein the oligosaccharide is linked to a functional group, optionally via a short spacer (see the section entitled Conjugation to a Non-Oligosaccharide Macromolecular Moiety). The *in vitro* glycosylation can be carried out in a suitable buffer at pH 4-7 in protein concentrations of 0.5-2 mg/ml and a volume of 0.02-2 ml. The activated mannose compound is present in 2-200 fold molar excess, and reactions are incubated at 4-25°C for periods of 0.1-3 hours. *In vitro* glycosylated GCB polypeptides are purified by dialysis and standard chromatographic techniques.

Other *in vitro* glycosylation methods are described, for example in WO 87/05330, by Aplin et al. (1981) *CRC Crit Rev. Biochem.* pp. 259-306, by Lundblad and Noyes *Chemical Reagents for Protein Modification* CRC Press Inc. Boca Raton, FL, by Yan and Wold (1984) *Biochemistry* 23: 3759-65, and by Doebber et al. (1982) *J. Biol. Chem.* 257: 2193-2199.

Furthermore, *in vitro* glycosylation to protein- and peptide-bound Gln-residues can be carried out by transglutaminases (TGases). Transglutaminases catalyse the transfer of donor amine-groups to protein- and peptide-bound Gln-residues in a so-called cross-linking reaction. The donor-amine groups can be protein- or peptide-bound e.g., as the  $\epsilon$ -amino-group in

Lys-residues or it can be part of a small or large organic molecule. An example of a small organic molecule functioning as amino-donor in TGase-catalysed cross-linking is putrescine (1,4-diaminobutane). An example of a larger organic molecule functioning as amino-donor in TGase-catalysed cross-linking is an amine-containing PEG (Sato et al. (1996) *Biochemistry* 35: 13072-13080).

TGases, in general, are highly specific enzymes, and not every Gln-residues exposed on the surface of a protein is accessible to TGase-catalysed cross-linking to amino-containing substances. In order to render a protein susceptible to TGase-catalysed cross-linking reactions stretches of amino acid sequence known to function very well as TGase substrates are inserted at convenient positions in the amino acid sequence encoding a GCB polypeptide. Several amino acid sequences are known to be or to contain excellent natural TGase substrates e.g., substance P, elafin, fibrinogen, fibronectin,  $\alpha_2$ -plasmin inhibitor,  $\alpha$ -caseins, and  $\beta$ -caseins and may thus be inserted into and thereby constitute part of the amino acid sequence of a polypeptide of the invention.

The nature and number of oligosaccharide moieties of a glycosylated polypeptide of the invention may be determined by a number of different methods known in the art e.g. by lectin binding studies (Reddy et al. (1985) *Biochem. Med.* 33: 200-210; Cummings (1994) *Meth. Enzymol.* 230: 66-86; *Protein Protocols* (Walker ed.) (1998) chapter 9); by reagent array analysis method (RAAM) sequencing of released oligosaccharides (Edge et al. (1992) *Proc. Natl. Acad. Sci. USA* 89: 6338-6342; Prime et al. (1996) *J. Chrom. A* 720: 263-274); by RAAM sequencing of released oligosaccharides in combination with mass spectrometry (Klausen et al. (1998) *Molecular Biotechnology* 9: 195-204); or by combining proteolytic degradation, glycopeptide purification by HPLC, exoglycosidase degradations and mass spectrometry (Krogh et al. (1997) *Eur. J. Biochem.* 244: 334-342). Specific methods for determining the glycosylation profile is described in the examples section hereinafter. Normally, the glycosylated polypeptide of the invention comprises 1-15 oligosaccharide moieties, such as 1-10 or 1-6 oligosaccharide moieties. Usually, at least one of these is attached to the peptide addition and further oligosaccharide structures are attached to the peptide addition or the polypeptide Pp.

Polypeptide of the invention conjugated to a second non-peptide moiety

It can be advantageous that the glycosylated polypeptide of the invention further comprises at least one second non-peptide moiety. The term "second non-peptide moiety" is

intended to indicate a non-peptide moiety different from an oligosaccharide moiety, e.g., a polymer molecule, a lipophilic compound and an organic derivatizing agent.

For this purpose the polypeptide must comprise at least one attachment group for the second non-peptide moiety. The attachment group can be one present on an amino acid residue, e.g., selected from the group consisting of the N-terminal or C-terminal amino acid residue of the polypeptide of the invention, lysine, cysteine, arginine, glutamine, aspartic acid, glutamic acid, serine, tyrosine, histidine, phenylalanine and tryptophan, or on an oligosaccharide moiety attached to the polypeptide. For instance, the attachment group for the non-peptide moiety is an epsilon-amino group.

It will be understood that an attachment group for the second non-peptide moiety may be provided by the N-terminal peptide addition, within the polypeptide Pp, and/or as a C-terminal peptide addition (having similar properties to those described above for the peptide addition X). In one embodiment, the peptide addition X comprising or contributing to an attachment site further comprises an attachment group for a second non-peptide moiety. For instance, the peptide addition may comprise 1-20, such as 1-10 attachment groups for a second non-peptide moiety. Such attachment groups may be distributed in a similar manner as that described immediately above for glycosylation sites. Also, the peptide addition X can comprise at least two attachment groups for the second non-peptide moiety.

Also, the polypeptide Pp can be a variant of a native polypeptide, which as compared to said native polypeptide, comprises at least one introduced and/or at least one removed attachment group for the second non-peptide moiety. For instance, the polypeptide Pp comprises at least one introduced attachment group, in particular 1-5 introduced attachment groups, such as 2-5 or 3-5 introduced attachment groups.

The attachment group is preferably located in a position that is exposed at the surface of the folded protein and thus accessible for conjugation to the polymer molecule. For instance, attachment to one or more polymer molecules increases the molecular weight of the polypeptide and can further serve to shield one or more epitopes thereof. The polymer molecule may be any of the molecules mentioned in the section entitled "Conjugation to a polymer molecule," but is preferably selected from the group consisting of linear or branched polyethylene glycol or polyalkylene oxide. Most preferably, the polymer molecule is mPEG-SPA, mPEG-SCM, mPEG-BTC from Shearwater Polymers, Inc, SC-PEG from Enzon, Inc.,

tresylated mPEG (US 5,880,255) or oxycarbonyl-oxy-N-dicarboxyimide PEG (US 5,122,614) (and the relevant attachment group is one present on a lysine or N-terminal residue).

Alternatively, the polymer molecule is an activated PEG molecule reactive with a cysteine residue, e.g., VS-PEG from Shearwater Polymers.

5           Especially, when the polypeptide Pp is an industrial enzyme, the second non-peptide moiety may be one which is capable of cross-linking and thereby of being immobilized on a suitable solid support. Such cross-linking polymers are available from Shearwater Polymers, Inc. It will be understood that the peptide addition of the polypeptide according to this embodiment comprises an attachment group for the cross-linking polymer in question. In  
10 connection with this embodiment, the polypeptide Pp is preferably an amyloglucosidase, an alpha-amylase, a glucose isomerase, an amidase, or a lipolytic enzyme.

In the following sections "Conjugation to a lipophilic compound," "Conjugation to a polymer molecule," and "Conjugation to an organic derivatizing agent" conjugation to specific types of non-peptide moieties is described.

15           It will be understood that a conjugation step of any method of the invention only finds relevance when a non-polypeptide moiety other than an *in vivo* attached oligosaccharide moiety is to be conjugated to the polypeptide, since *in vivo* glycosylation takes place during the expression step when using an appropriate glycosylating host cell as expression host. Accordingly, whenever a conjugation step occurs in the present invention this is intended to be  
20 conjugation to a non-polypeptide moiety other than an oligosaccharide moiety attached by *in vivo* glycosylation during expression in a glycosylating organism. *In vitro* glycosylation methods are described in the section entitled "glycosylation."

#### Conjugation to a lipophilic compound

25           The polypeptide and the lipophilic compound can be conjugated to each other, either directly or by use of a linker. The lipophilic compound can be a natural compound such as a saturated or unsaturated fatty acid, a fatty acid diketone, a terpene, a prostaglandin, a vitamine, a carotenoide or steroide, or a synthetic compound such as a carbon acid, an alcohol, an amine and sulphonic acid with one or more alkyl-, aryl-, alkenyl- or other multiple unsaturated  
30 compounds. Furthermore, the lipophilic compound may be any of the lipophilic substituents disclosed in WO 97/31022, the contents of which are incorporated herein by reference. The conjugation between the polypeptide and the lipophilic compound, optionally through a linker

can be done according to methods known in the art, e.g., as described by Bodanszky (1976), in *Peptide Synthesis*, John Wiley, New York and in WO 96/12505 and further as described in WO 97/31022.

#### Conjugation to a polymer molecule

5 The polymer molecule to be coupled to the polypeptide of the invention can be any suitable polymer molecule, such as a natural or synthetic homo-polymer or heteropolymer, typically with a molecular weight in the range of 300-100,000 Da, such as 300-20,000 Da, more preferably in the range of 500-10,000 Da, even more preferably in the range of 500-5000 Da.

10 Examples of homo-polymers include a polyol (i.e., poly-OH), a polyamine (i.e., poly-NH<sub>2</sub>) and a polycarboxylic acid (i.e., poly-COOH). A hetero-polymer is a polymer that comprises different coupling groups, such as a hydroxyl group and an amine group.

15 Examples of suitable polymer molecules include polymer molecules selected from the group consisting of polyalkylene oxide (PAO), including polyalkylene glycol (PAG), such as polyethylene glycol (PEG) and polypropylene glycol (PPG), branched PEGs, poly-vinyl alcohol (PVA), poly-carboxylate, poly-(vinylpyrrolidone), polyethylene-co-maleic acid anhydride, polystyrene-co-malic acid anhydride, dextran, including carboxymethyl-dextran, or any other biopolymer suitable for the intended purpose, such as for reducing immunogenicity and/or increasing functional *in vivo* half-life and/or serum half-life, or for providing immobilization properties to the polypeptide (as discussed in the section entitled "Polypeptide of interest." Another example of a polymer molecule is human albumin or another abundant plasma protein. Generally, polyalkylene glycol-derived polymers are biocompatible, non-toxic, non-antigenic, non-immunogenic, have various water solubility properties, and are easily excreted from living organisms.

20 PEG is the preferred polymer molecule for reducing immunogenicity, 25 allergenicity and/or increasing half-life, since it has only few reactive groups capable of cross-linking compared, e.g., to polysaccharides such as dextran, and the like. In particular, monofunctional PEG, e.g., methoxypolyethylene glycol (mPEG), is of interest since its coupling chemistry is relatively simple (only one reactive group is available for conjugating with attachment groups on the polypeptide). Consequently, the risk of cross-linking is eliminated, the 30 resulting polypeptide conjugates are more homogeneous and the reaction of the polymer molecules with the polypeptide is easier to control.

To effect covalent attachment of the polymer molecule(s) to the polypeptide, the hydroxyl end groups of the polymer molecule must be provided in activated form, i.e., with reactive functional groups. Suitable activated polymer molecules are commercially available, e.g., from Shearwater Polymers, Inc., Huntsville, AL, USA. Alternatively, the polymer molecules can be activated by conventional methods known in the art, e.g., as disclosed in WO 90/13540. Specific examples of activated linear or branched polymer molecules for use in the present invention are described in the Shearwater Polymers, Inc. 1997 and 2000 Catalogs (Functionalized Biocompatible Polymers for Research and pharmaceuticals, Polyethylene Glycol and Derivatives, incorporated herein by reference). Specific examples of activated PEG polymers include the following linear PEGs: NHS-PEG (e.g., SPA-PEG, SSPA-PEG, SBA-PEG, SS-PEG, SSA-PEG, SC-PEG, SG-PEG, and SCM-PEG), and NOR-PEG), BTC-PEG, EPOX-PEG, NCO-PEG, NPC-PEG, CDI-PEG, ALD-PEG, TRES-PEG, VS-PEG, IODO-PEG, and MAL-PEG, and branched PEGs such as PEG2-NHS and those disclosed in US 5,932,462 and US 5,643,575, both of which are incorporated herein by reference. Furthermore, the following publications, incorporated herein by reference, disclose useful polymer molecules and/or PEGylation chemistries: US 5,824,778, US 5,476,653, WO 97/32607, EP 229,108, EP 402,378, US 4,902,502, US 5,281,698, US 5,122,614, US 5,219,564, WO 92/16555, WO 94/04193, WO 94/14758, WO 94/17039, WO 94/18247, WO 94/28024, WO 95/00162, WO 95/11924, WO95/13090, WO 95/33490, WO 96/00080, WO 97/18832, WO 98/41562, WO 98/48837, WO 99/32134, WO 99/32139, WO 99/32140, WO 96/40791, WO 98/32466, WO 95/06058, EP 439 508, WO 97/03106, WO 96/21469, WO 95/13312, EP 921 131, US 5,736,625, WO 98/05363, EP 809 996, US 5,629,384, WO 96/41813, WO 96/07670, US 5,473,034, US 5,516,673, EP 605 963, US 5,382,657, EP 510 356, EP 400 472, EP 183 503 and EP 154 316.

The conjugation of the polypeptide and the activated polymer molecules is conducted by use of any conventional method, e.g., as described in the following references (which also describe suitable methods for activation of polymer molecules): R.F. Taylor (1991) *Protein immobilisation: Fundamental and applications* Marcel Dekker, N.Y.; S.S. Wong (1992) *Chemistry of Protein Conjugation and Crosslinking* CRC Press, Boca Raton; G.T. Hermanson et al. (1993) *Immobilized Affinity Ligand Techniques* Academic Press, N.Y.). The skilled person will be aware that the activation method and/or conjugation chemistry to be used depends on the attachment group(s) of the polypeptide (examples of which are given further above), as well as

the functional groups of the polymer (e.g., being amine, hydroxyl, carboxyl, aldehyde, sulfhydryl, succinimidyl, maleimide, vinylsulfone or haloacetate). The PEGylation can be directed towards conjugation to all available attachment groups on the polypeptide (i.e., such attachment groups that are exposed at the surface of the polypeptide) or can be directed towards one or more specific attachment groups, e.g., the N-terminal amino group (US 5,985,265). Furthermore, the conjugation can be achieved in one step or in a stepwise manner (e.g., as described in WO 99/55377).

It will be understood that the PEGylation is designed so as to produce the optimal molecule with respect to the number of PEG molecules attached, the size and form of such molecules (e.g., whether they are linear or branched), and where in the polypeptide such molecules are attached. For instance, the molecular weight of the polymer to be used can be chosen on the basis of the desired effect to be achieved. For instance, if the primary purpose of the conjugation is to achieve a polypeptide having a high molecular weight (e.g., to reduce renal clearance) it is usually desirable to conjugate as few high MW polymer molecules as possible to obtain the desired molecular weight. When a high degree of epitope shielding is desirable this can be obtained by use of a sufficiently high number of low molecular weight polymer molecules (e.g., with a molecular weight of about 5,000 Da) to effectively shield all or most epitopes of the polypeptide. For instance, 2-8, such as 3-6 such polymers can be used.

In connection with conjugation to only a single attachment group on the protein (as described in US 5,985,265), it can be advantageous that the polymer molecule, which can be linear or branched, has a high molecular weight, e.g., about 20 kDa.

Normally, the polymer conjugation is performed under conditions aiming at reacting all available polymer attachment groups with polymer molecules. Typically, the molar ratio of activated polymer molecules to polypeptide is up to about 1000-1, in particular 200-1, preferably 100-1, such as 10-1 or 5-1, but also equimolar ratios can be used in order to obtain optimal reaction.

It is also contemplated according to the invention to couple the polymer molecules to the polypeptide through a linker. Suitable linkers are well known to the skilled person. A preferred example is cyanuric chloride (Abuchowski et al. (1977) *J. Biol. Chem.* 252: 3578-3581; US 4,179,337; Shafer et al. (1986) *J. Polym. Sci. Polym. Chem. Ed.* 24: 375-378).

Subsequent to the conjugation residual activated polymer molecules are blocked according to methods known in the art, e.g., by addition of primary amine to the reaction mixture, and the resulting inactivated polymer molecules are removed by a suitable method.

In a specific embodiment, the polypeptide of the invention is one that comprises one or more PEG molecules attached to the peptide addition, but not to the polypeptide P. For instance, the PEG molecule is attached to one or more cysteine residues present in the peptide addition X and, if necessary, one or more cysteine residues have been removed from the polypeptide P of interest in order to avoid conjugation thereto.

In another specific embodiment, the polypeptide of the invention comprises at least one PEG molecule attached to a lysine residue of the peptide addition X, in particular a linear or branched PEG molecule with a molecular weight of at least 5kDa.

#### Methods of preparing a polypeptide of the invention

The invention further comprises a method of producing the polypeptide of the invention, which method comprises culturing a host cell transformed or transfected with a nucleotide sequence encoding the polypeptide under conditions permitting the expression of the polypeptide, and recovering the polypeptide from the culture.

Apart from recombinant production, polypeptides of the invention may be produced, albeit less efficiently, by chemical synthesis or a combination of chemical synthesis and recombinant DNA technology.

The nucleotide sequence of the invention encoding a polypeptide of the invention may be constructed by isolating or synthesizing a nucleotide sequence encoding the parent polypeptide and fusing a nucleotide sequence encoding the relevant peptide addition in accordance with established technologies. To the extent amino acid modifications are to be made in the parent polypeptide, these are conveniently done by mutagenesis, e.g., using site-directed mutagenesis in accordance with well-known methods, e.g., as described in Nelson and Long (1989) *Analytical Biochemistry* 180: 147-151, random mutagenesis, or shuffling.

The nucleotide sequence may be prepared by chemical synthesis, e.g., by using an oligonucleotide synthesizer, wherein oligonucleotides are designed based on the amino acid sequence of the desired polypeptide, and preferably selecting those codons that are favored in the host cell in which the recombinant polypeptide will be produced. For example, several small oligonucleotides coding for portions of the desired polypeptide may be synthesized and

assembled by polymerase chain reaction (PCR), ligation or ligation chain reaction (LCR). The individual oligonucleotides typically contain 5' or 3' overhangs for complementary assembly.

Once assembled (by synthesis, site-directed mutagenesis or another method), the nucleotide sequence encoding the polypeptide may be inserted into a recombinant vector and operably linked to control sequences necessary for expression of thereof in the desired transformed host cell.

It should of course be understood that not all vectors and expression control sequences function equally well to express the nucleotide sequence encoding the polypeptide part of the invention. Neither will all hosts function equally well with the same expression system. However, one of skill in the art can make a selection among these vectors, expression control sequences and hosts without undue experimentation. For example, in selecting a vector, the host must be considered because the vector must replicate in it or be able to integrate into the chromosome. The vector's copy number, the ability to control that copy number, and the expression of any other proteins encoded by the vector, such as antibiotic markers, should also be considered. In selecting an expression control sequence, a variety of factors should also be considered. These include, for example, the relative strength of the sequence, its controllability, and its compatibility with the nucleotide sequence encoding the polypeptide, particularly as regards potential secondary structures. Hosts should be selected by consideration of their compatibility with the chosen vector, the toxicity of the product coded for by the nucleotide sequence, their secretion characteristics, their ability to fold the polypeptide correctly, their fermentation or culture requirements, and the ease of purification of the products coded for by the nucleotide sequence.

The recombinant vector may be an autonomously replicating vector, i.e., a vector existing as an extrachromosomal entity, the replication of which is independent of chromosomal replication, e.g., a plasmid. Alternatively, the vector is one which, when introduced into a host cell, is integrated into the host cell genome and replicated together with the chromosome(s) into which it has been integrated.

The vector is preferably an expression vector, in which the nucleotide sequence encoding the polypeptide of the invention is operably linked to additional segments required for transcription of the nucleotide sequence. The vector is typically derived from plasmid or viral DNA. A number of suitable expression vectors for expression in the host cells mentioned herein

are commercially available or described in the literature. Useful expression vectors for eukaryotic hosts, include, for example, vectors comprising expression control sequences from SV40, bovine papilloma virus, adenovirus and cytomegalovirus. Specific vectors are, e.g., pCDNA3.1(+)\Hyg (Invitrogen, Carlsbad, CA, USA) and pCI-neo (Stratagene, La Jolla, CA, USA). Useful expression vectors for yeast cells include the 2 $\mu$  plasmid and derivatives thereof, the POT1 vector (US 4,931,373), the pJSO37 vector described in (Okkels, Ann. New York Acad. Sci. 782, 202-207, 1996) and pPICZ A, B or C (Invitrogen, Carlsbad, CA, USA). Useful vectors for insect cells include pVL941, pBG311 (Cate et al. (1986) "Isolation of the Bovine and Human Genes for Mullerian Inhibiting Substance And Expression of the Human Gene In Animal Cells" *Cell* 45: 685-98, pBluebac 4.5 and pMelbac (both available from Invitrogen, Carlsbad, CA, USA).

Other vectors for use in this invention include those that allow the nucleotide sequence encoding the polypeptide of the invention to be amplified in copy number. Such amplifiable vectors are well known in the art. They include, for example, vectors able to be amplified by DHFR amplification (see, e.g., Kaufman, U.S. Pat. No. 4,470,461, Kaufman and Sharp (1982) "Construction Of A Modular Dihydrofolate Reductase cDNA Gene: Analysis Of Signals Utilized For Efficient Expression" *Mol. Cell. Biol.* 2: 1304-19) and glutamine synthetase ("GS") amplification (see, e.g., US 5,122,464 and EP 338,841).

The recombinant vector may further comprise a DNA sequence enabling the vector to replicate in the host cell in question. An example of such a sequence (when the host cell is a mammalian cell) is the SV40 origin of replication. When the host cell is a yeast cell, suitable sequences enabling the vector to replicate are the yeast plasmid 2 $\mu$  replication genes REP 1-3 and origin of replication.

The vector may also comprise a selectable marker, e.g., a gene the product of which complements a defect in the host cell, such as the gene coding for dihydrofolate reductase (DHFR) or the *Schizosaccharomyces pombe* TPI gene (described by P.R. Russell (1985) *Gene* 40: 125-130), or one which confers resistance to a drug, e.g., ampicillin, kanamycin, tetracyclin, chloramphenicol, neomycin, hygromycin or methotrexate. For filamentous fungi, selectable markers include amdS, pyrG, arcB, niaD, sC.

The term "control sequences" is defined herein to include all components, which are necessary or advantageous for the expression of the polypeptide of the invention. Each

control sequence may be native or foreign to the nucleic acid sequence encoding the polypeptide. Such control sequences include, but are not limited to, a leader, polyadenylation sequence, propeptide sequence, promoter, enhancer or upstream activating sequence, signal peptide sequence, and transcription terminator. At a minimum, the control sequences include a promoter operably linked to the nucleotide sequence encoding the polypeptide.

“Operably linked” refers to the covalent joining of two or more nucleotide sequences, by means of enzymatic ligation or otherwise, in a configuration relative to one another such that the normal function of the sequences can be performed. For example, the nucleotide sequence encoding a presequence or secretory leader is operably linked to a nucleotide sequence for a polypeptide if it is expressed as a preprotein that participates in the secretion of the polypeptide: a promoter or enhancer is operably linked to a coding sequence if it affects the transcription of the sequence; a ribosome binding site is operably linked to a coding sequence if it is positioned so as to facilitate translation. Generally, “operably linked” means that the nucleotide sequences being linked are contiguous and, in the case of a secretory leader, contiguous and in reading phase. Linking is accomplished by ligation at convenient restriction sites. If such sites do not exist, then synthetic oligonucleotide adaptors or linkers are used, in conjunction with standard recombinant DNA methods.

A wide variety of expression control sequences may be used in the present invention. Such useful expression control sequences include the expression control sequences associated with structural genes of the foregoing expression vectors as well as any sequence known to control the expression of genes of prokaryotic or eukaryotic cells or their viruses, and various combinations thereof.

Examples of suitable control sequences for directing transcription in mammalian cells include the early and late promoters of SV40 and adenovirus, e.g., the adenovirus 2 major late promoter, the MT-1 (metallothionein gene) promoter, the human cytomegalovirus immediate-early gene promoter (CMV), the human elongation factor 1 $\alpha$  (EF-1 $\alpha$ ) promoter, the *Drosophila* minimal heat shock protein 70 promoter, the Rous Sarcoma Virus (RSV) promoter, the human ubiquitin C (UbC) promoter, the human growth hormone terminator, SV40 or adenovirus Elb region polyadenylation signals and the Kozak consensus sequence (Kozak (1987) *J Mol Biol* 196: 947-50).

In order to improve expression in mammalian cells a synthetic intron may be inserted in the 5' untranslated region of the nucleotide sequence encoding the polypeptide of the invention. An example of a synthetic intron is the synthetic intron from the plasmid pCI-Neo (available from Promega Corporation, WI, USA).

5           Examples of suitable control sequences for directing transcription in insect cells include the polyhedrin promoter, the P10 promoter, the *Autographa californica* polyhedrosis virus basic protein promoter, the baculovirus immediate early gene 1 promoter and the baculovirus 39K delayed-early gene promoter, and the SV40 polyadenylation sequence.

10           Examples of suitable control sequences for use in yeast host cells include the promoters of the yeast  $\alpha$ -mating system, the yeast triose phosphate isomerase (TPI) promoter, promoters from yeast glycolytic genes or alcohol dehydrogenase genes, the ADH2-4c promoter and the inducible GAL promoter.

15           Examples of suitable control sequences for use in filamentous fungal host cells include the ADH3 promoter and terminator, a promoter derived from the genes encoding *Aspergillus oryzae* TKA amylase triose phosphate isomerase or alkaline protease, an *A. niger*  $\alpha$ -amylase, *A. niger* or *A. nidulans* glucoamylase, *A. nidulans* acetamidase, *Rhizomucor miehei* aspartic proteinase or lipase, the TPI1 terminator and the ADH3 terminator.

20           The nucleotide sequence of the invention may or may not also include a nucleotide sequence that encode a signal peptide. The signal peptide is present when the polypeptide is to be secreted from the cells in which it is expressed. Such signal peptide, if present, should be one recognized by the cell chosen for expression of the polypeptide. The signal peptide may be homologous (e.g., be that normally associated with the parent polypeptide in question) or heterologous (i.e., originating from another source than the parent polypeptide) to the polypeptide or may be homologous or heterologous to the host cell, i.e., be a signal peptide normally expressed from the host cell or one which is not normally expressed from the host cell. Accordingly, the signal peptide may be prokaryotic, e.g., derived from a bacterium, or eukaryotic, e.g., derived from a mammalian, or insect, filamentous fungal or yeast cell.

25           The presence or absence of a signal peptide will, e.g., depend on the expression host cell used for the production of the polypeptide, the protein to be expressed (whether it is an intracellular or extracellular protein) and whether it is desirable to obtain secretion. For use in filamentous fungi, the signal peptide may conveniently be derived from a gene encoding an

*Aspergillus* sp. amylase or glucoamylase, a gene encoding a *Rhizomucor miehei* lipase or protease or a *Humicola lanuginosa* lipase. The signal peptide is preferably derived from a gene encoding *A. oryzae* TAKA amylase, *A. niger* neutral  $\alpha$ -amylase, *A. niger* acid-stable amylase, or *A. niger* glucoamylase. For use in insect cells, the signal peptide may conveniently be derived from an insect gene (cf. WO 90/05783), such as the lepidopteran *Manduca sexta* adipokinetic hormone precursor, (cf. US 5,023,328), the honeybee melittin (Invitrogen, Carlsbad, CA, USA), ecdysteroid UDPglucosyltransferase (egt) (Murphy et al. (1993) *Protein Expression and Purification* 4: 349-357, or human pancreatic lipase (hpl) (*Methods in Enzymology* (1997) 284: 262-272).

Specific examples of signal peptides for use in mammalian cells include that of human glucocerebrosidase apparent from the examples hereinafter or the murine Ig kappa light chain signal peptide (Coloma, M (1992) *J. Imm. Methods* 152: 89-104). For use in yeast cells suitable signal peptides have been found to be the  $\alpha$ -factor signal peptide from *S. cerevisiae*. (cf. US 4,870,008), the signal peptide of mouse salivary amylase (cf. O. Hagenbuchle et al. (1981) *Nature* 289: 643-646), a modified carboxypeptidase signal peptide (cf. L.A. Valls et al.(1987) *Cell* 48: 887-897), the yeast BAR1 signal peptide (cf. WO 87/02670), and the yeast aspartic protease 3 (YAP3) signal peptide (cf. M. Egel-Mitani et al. (1990) *Yeast* 6: 127-137).

Any suitable host may be used to produce the polypeptide of the invention, including bacteria, fungi (including yeasts), plant, insect, mammal, or other appropriate animal cells or cell lines, as well as transgenic animals or plants. When a non-glycosylating organism such as *E. coli* is used, and the polypeptide is to be a glycosylated polypeptide, the expression in *E. coli* is preferably followed by suitable *in vitro* glycosylation.

Examples of bacterial host cells include grampositive bacteria such as strains of *Bacillus*, e.g., *B. brevis* or *B. subtilis*, *Pseudomonas* or *Streptomyces*, or gramnegative bacteria, such as strains of *E. coli*. The introduction of a vector into a bacterial host cell may, for instance, be effected by protoplast transformation (see, e.g., Chang and Cohen (1979) *Molecular General Genetics* 168: 111-115), using competent cells (see, e.g., Young and Spizizin (1961) *Journal of Bacteriology* 81: 823-829, or Dubnau and Davidoff-Abelson (1971) *Journal of Molecular Biology* 56: 209-221), electroporation (see, e.g., Shigekawa and Dower (1988) *Biotechniques* 6: 742-751), or conjugation (see, e.g., Koehler and Thorne (1987) *Journal of Bacteriology* 169: 5771-5278).

Examples of suitable filamentous fungal host cells include strains of *Aspergillus*, e.g., *A. oryzae*, *A. niger*, or *A. nidulans*, *Fusarium* or *Trichoderma*. Fungal cells may be transformed by a process involving protoplast formation, transformation of the protoplasts, and regeneration of the cell wall in a manner known *per se*. Suitable procedures for transformation of *Aspergillus* host cells are described in EP 238 023 and US 5,679,543. Suitable methods for transforming *Fusarium* species are described by Malardier et al. (1989) *Gene* 78: 147-156 and WO 96/00787. Yeast may be transformed using the procedures described by Becker and Guarente, In Abelson, J.N. and Simon, M.I., editors, *Guide to Yeast Genetics and Molecular Biology, Methods in Enzymology*, Volume 194, pp 182-187, Academic Press, Inc., New York; Ito et al. (1983) *Journal of Bacteriology* 153: 163; and Hinnen et al. (1978) *Proceedings of the National Academy of Sciences USA* 75: 1920.

When the polypeptide of the invention is to be *in vivo* glycosylated, the host cell is selected from a group of host cells capable of generating the desired glycosylation of the polypeptide. Thus, the host cell may advantageously be selected from a yeast cell, insect cell, or mammalian cell.

Examples of suitable yeast host cells include strains of *Saccharomyces*, e.g., *S. cerevisiae*, *Schizosaccharomyces*, *Kluyveromyces*, *Pichia*, such as *P. pastoris* or *P. methanolica*, *Hansenula*, such as *H. polymorpha* or *glabrata*. Of particular interest are yeast glycosylation mutant cells, e.g., derived from *S. cerevisiae*, *P. pastoris* or *Hansenula* spp. (e.g., the *S. cerevisiae* glycosylation mutants *och1*, *och1 mnm1* or *och1 mnm1 alg3* described by Nagasu et al. (1992) *Yeast* 8: 535-547 and Nakanisho-Shindo et al. (1993) *J. Biol. Chem.* 268: 26338-26345. Methods for transforming yeast cells with heterologous DNA and producing heterologous polypeptides therefrom are disclosed by Clontech Laboratories, Inc, Palo Alto, CA, USA (in the product protocol for the Yeastmaker™ Yeast Transformation System Kit), and by Reeves et al. (1992) *FEMS Microbiology Letters* 99: 193-198, Manivasakam and Schiestl (1993) *Nucleic Acids Research* 21: 4414-4415 and Ganeva et al. (1994) *FEMS Microbiology Letters* 121: 159-164.

Examples of suitable insect host cells include a *Lepidoptera* cell line, such as *Spodoptera frugiperda* (Sf9 or Sf21) or *Trichoplusia ni* cells (High Five) (US 5,077,214). Transformation of insect cells and production of heterologous polypeptides therein may be performed as described by Invitrogen, Carlsbad, CA, USA.

Examples of suitable mammalian host cells include Chinese hamster ovary (CHO) cell lines, (e.g., CHO-K1; ATCC CCL-61), Green Monkey cell lines (COS) (e.g., COS 1 (ATCC CRL-1650), COS 7 (ATCC CRL-1651)); mouse cells (e.g., NS/O), Baby Hamster Kidney (BHK) cell lines (e.g., ATCC CRL-1632 or ATCC CCL-10), and human cells (e.g., HEK 293 (ATCC CRL-1573)), as well as plant cells in tissue culture. Additional suitable cell lines are known in the art and available from public depositories such as the American Type Culture Collection, Rockville, Maryland. Of interest for the present purpose are a mammalian glycosylation mutant cell line, such as CHO-LEC1, CHOL-LEC2 or CHO-LEC18 (CHO-LEC1: Stanley et al. (1975) *Proc. Natl. Acad. USA* 72: 3323-3327 and Grossmann et al. (1995) *J. Biol. Chem.* 270: 29378-29385, CHO-LEC18: Raju et al. (1995) *J. Biol. Chem.* 270: 30294-30302).

Methods for introducing exogenous DNA into mammalian host cells include calcium phosphate-mediated transfection, electroporation, DEAE-dextran mediated transfection, liposome-mediated transfection, viral vectors and the transfection method described by Life Technologies Ltd, Paisley, UK using Lipofectamin 2000. These methods are well known in the art and e.g., described by Ausbel et al. (eds.) (1996) *Current Protocols in Molecular Biology* John Wiley & Sons, New York, USA. The cultivation of mammalian cells are conducted according to established methods, e.g., as disclosed in Jenkins, Ed. (1999) *Animal Cell Biotechnology, Methods and Protocols* Human Press Inc, Totowa, New Jersey, USA; and Harrison and Rae (1997) *General Techniques of Cell Culture* Cambridge University Press.

In the production methods of the present invention, cells are cultivated in a nutrient medium suitable for production of the polypeptide using methods known in the art. For example, cells are cultivated by shake flask cultivation, small-scale or large-scale fermentation (including continuous, batch, fed-batch, or solid state fermentations) in laboratory or industrial fermenters performed in a suitable medium and under conditions allowing the polypeptide to be expressed and/or isolated. The cultivation takes place in a suitable nutrient medium comprising carbon and nitrogen sources and inorganic salts, using procedures known in the art. Suitable media are available from commercial suppliers or may be prepared according to published compositions (e.g., in catalogues of the American Type Culture Collection). If the polypeptide is secreted into the nutrient medium, the polypeptide can be recovered directly from the medium. If the polypeptide is not secreted, it can be recovered from cell lysates.

The resulting polypeptide may be recovered by methods known in the art. For example, the polypeptide may be recovered from the nutrient medium by conventional procedures including, but not limited to, centrifugation, filtration, extraction, spray drying, evaporation, or precipitation.

5 The polypeptides may be purified by a variety of procedures known in the art including, but not limited to, chromatography (*e.g.*, ion exchange, affinity, hydrophobic, chromatofocusing, and size exclusion), electrophoretic procedures (*e.g.*, preparative isoelectric focusing), differential solubility (*e.g.*, ammonium sulfate precipitation), SDS-PAGE, or extraction (see, *e.g.*, J-C Janson and Lars Ryden, editors (1989) *Protein Purification* VCH  
10 Publishers, New York,).

#### Other methods of the invention

In accordance with a specific aspect a nucleotide sequence encoding the polypeptide of the invention is prepared by a method comprising:

- 15 a) subjecting a nucleotide sequence encoding the polypeptide Pp to elongation mutagenesis;  
b) expressing the mutated nucleotide sequence obtained in step a) in a suitable host cell;  
c) optionally conjugating polypeptides expressed in step b) to a second non-peptide moiety;  
d) selecting polypeptides of step b) or c) which comprises at least one oligosaccharide moiety and optionally second non-peptide moiety attached to the peptide addition part of the polypeptide; and,  
20 e) isolating a nucleotide sequence encoding the polypeptide selected in step d).

In the present context the term “elongation mutagenesis” is intended to indicate any manner in which the nucleotide sequence encoding the parent polypeptide Pp can be extended to further encode the peptide addition. For instance, a nucleotide sequence encoding a peptide addition of a suitable length may be synthesized and fused to a nucleotide sequence  
25 encoding the polypeptide Pp. The resulting fused nucleotide sequence may then be subjected to further modification by any suitable method, *e.g.*, one which involves gene shuffling, other recombination between nucleotide sequences, random mutagenesis, random elongation mutagenesis or any combination of these methods. Such methods are further described in the Methods section herein.

The expression and optional conjugation steps are conducted as described in further detail elsewhere in the present application, and the selection step d) using any suitable method available in the art.

5 In one embodiment, the above method further comprises screening polypeptides resulting from step b) or c) for at least one improved property, in particular any of those improved properties listed herein, prior to the selection step, and wherein the selection step d) further comprises selecting polypeptides having such improved property.

10 Furthermore, in the above method the elongation mutagenesis can be conducted so as to enrich for codons encoding a glycosylation site and/or an amino acid residue comprising an attachment group for a second non-peptide moiety., in particular an *in vivo* glycosylation site.

Still further, the above method can comprise subjecting the part of the nucleotide sequence encoding the polypeptide Pp of interest to mutagenesis to remove and/or introduce glycosylation site(s) and/or amino acid residue(s) comprising an attachment group for the second non-peptide moiety. The nucleotide sequence may be subjected to any type of mutagenesis, e.g.,  
15 any of those described herein. The mutagenesis of the nucleotide sequence encoding the polypeptide Pp of interest can be conducted prior to assembling the sequence with that encoding the peptide addition, concomitantly with or after any mutagenesis of the peptide addition part of the assembled nucleotide sequence.

20 In a further aspect, the invention relates to a method of producing a glycosylated polypeptide encoded by a nucleotide sequence of the invention prepared by the above method, wherein the nucleotide sequence encoding the polypeptide selected in step c) is expressed in a glycosylating host cell and the resulting glycosylated expressed polypeptide is recovered.

In a still further aspect the invention relates to a method of improving one or more selected properties of a polypeptide Pp of interest, which method comprises:

- 25 a) preparing a nucleotide sequence encoding a polypeptide comprising or consisting essentially of the primary structure  $\text{NH}_2\text{-X-Pp-COOH}$ , wherein X is a peptide addition comprising or contributing to a glycosylation site and/or an attachment group for a second non-peptide moiety that is capable of conferring the selected improved property/ies to the polypeptide Pp;  
b) expressing the nucleotide sequence of a) in an suitable host cell;  
30 c) optionally conjugating the expressed polypeptide of b) to a second non-peptide moiety; and,  
d) recovering the polypeptide resulting from step b) or c).

For instance, the polypeptide is any of those described herein. For instance the nucleotide sequence of step a) is prepared by subjecting a nucleotide sequence encoding the polypeptide Pp to elongation mutagenesis, e.g., to enrich for codons encoding an amino acid residue comprising or contributing to a glycosylation site and/or an attachment group for a second non-peptide moiety, in particular an *in vivo* glycosylation site. Also, in the preparation of the nucleotide sequence of a), the part of the nucleotide sequence encoding the polypeptide Pp can be subjected to mutagenesis to remove and/or introduce glycosylation site(s) and/or attachment group(s) for a second non-peptide moiety.

The method according to this aspect can further comprise a screening step (after step c)), wherein the polypeptide resulting from step b) or c) is screened for one or more improved properties, in particular any of those improved properties which are described hereinabove.

Usually, when a polypeptide has been selected in a screening step of a method of the invention the nucleotide sequence encoding the polypeptide is isolated and used for expression of larger amounts of the polypeptide. The amino acid sequence of the resulting polypeptide is determined and the polypeptide may be subjected to conjugation in a larger scale. Subsequently, the polypeptide is assayed with respect to the property to be improved.

#### Uses of a polypeptide of the invention

It will be understood that polypeptides of the invention can be used for a variety of purposes, depending on the type and nature of polypeptide. For instance, it is contemplated that a polypeptide of the invention prepared from a therapeutic polypeptide is useful for the same therapeutic purposes as the parent polypeptide, i.e., for the treatment of a particular disease. Accordingly, the polypeptide of the invention may be formulated into a pharmaceutical composition. Also, when the polypeptide of the invention is an *in vivo* glycosylated polypeptide which does not comprise any other type of non-peptide moiety, a nucleotide sequence encoding the polypeptide can be used in gene therapy in accordance with established principles. When the polypeptide Pp is an antigen the polypeptide of the invention may be provided in the form of a vaccine.

## METHODS

### NUCLEOTIDE SEQUENCE MODIFICATION METHODS

For example, a peptide addition may be constructed from two or more nucleotide sequences encoding a polypeptide of interest with a peptide addition, the sequences being sufficiently homologous to allow recombination between the sequences, in particular in the part thereof encoding the peptide addition. The combination of nucleotide sequences or sequence parts is conveniently conducted by methods known in the art, for instance methods which involve homologous cross-over such as disclosed in US 5,093,257, or methods which involve gene shuffling, i.e., recombination between two or more homologous nucleotide sequences resulting in new nucleotide sequences having a number of nucleotide alterations when compared to the starting nucleotide sequences. In order for homology based nucleic acid shuffling to take place the relevant parts of the nucleotide sequences are preferably at least 50% identical, such as at least 60% identical, more preferably at least 70% identical, such as at least 80% identical. The recombination can be performed in vitro or in vivo. Examples of suitable in vitro gene shuffling methods are disclosed by Stemmer et al (1994) *Proc. Natl. Acad. Sci. USA* 91: 10747-10751; Stemmer (1994) *Nature* 370: 389-391; Smith (1994) *Nature* 370: 324-325; Zhao et al. *Nat. Biotechnol.* (1998) 16(3): 258-61; Zhao H. and Arnold, *Nucleic Acids Research* (1997) 25: 1307-1308; Shao et al. (1998) *Nucleic Acids Research* 26(2): 681-83; and WO 95/17413. Example of a suitable in vivo shuffling method is disclosed in WO 97/07205.

Furthermore, a peptide addition can be constructed by preparing a randomly mutagenized library, conveniently prepared by subjecting a nucleotide sequence encoding the polypeptide of the invention or the peptide addition to random mutagenesis to create a large number of mutated nucleotide sequences. While the random mutagenesis can be entirely random, both with respect to where in the nucleotide sequence the mutagenesis occurs and with respect to the nature of mutagenesis, it is preferably conducted so as to randomly mutate only the part of the sequence that encode the peptide addition. Also, the random mutagenesis can be directed towards introducing certain types of amino acid residues, in particular amino acid residues containing an attachment group, at random into the polypeptide molecule or at random into peptide addition part thereof. Besides substitutions, random mutagenesis can also cover

random introduction of insertions or deletions. Preferably, the insertions are made in reading frame, e.g., by performing multiple introduction of three nucleotides as described by Hallet et al. (1997) *Nucleic Acids Res* 25(9): 1866-7 and Sondek and Shrotle (1992) *Proc Natl. Acad. Sci USA* 89(8): 3581-5.

5                   The random mutagenesis (either of the whole nucleotide sequence or more preferably the part thereof encoding the peptide addition) can be performed by any suitable method. For example, the random mutagenesis is performed using a suitable physical or chemical mutagenizing agent, a suitable oligonucleotide, PCR generated mutagenesis or any combination of these mutagenizing agents and/or other methods according to state of the art  
10                   technology, e.g., as disclosed in WO 97/07202.

                    Error prone PCR generated mutagenesis, e.g., as described by J.O. Deshler (1992) *GATA* 9(4): 103-106 and Leung et al. (1989) *Technique* 1: 11-15, is particularly useful for mutagenesis of longer peptide stretches (corresponding to nucleotide sequences containing more than 100 bp) or entire genes, and are preferably performed under conditions that increase the  
15                   misincorporation of nucleotides.

                    Random mutagenesis based on doped or spiked oligonucleotides or by specific sequence oligonucleotides, is of particular use for mutagenesis of the part of the nucleotide sequence encoding the peptide addition.

                    Random mutagenesis of the part of the nucleotide sequence encoding the peptide  
20                   addition can be performed using PCR generated mutagenesis, in which one or more suitable oligonucleotide primers flanking the area to be mutagenized are used. In addition, doping or spiking with oligonucleotides can be used to introduce mutations so as to remove or introduce attachment groups for the relevant non-peptide moiety. State of the art knowledge and computer programs (e.g., as described by Siderovski DP and Mak TW (1993) *Comput. Biol. Med.* 23: 463-  
25                   474 and Jensen et al. (1998) *Nucleic Acids Research* 26, No. 3) can be used for calculating the most optimal nucleotide mixture for a given amino acid preference. The oligonucleotides can be incorporated into the nucleotide sequence encoding the peptide addition by any published technique using e.g., PCR, LCR or any DNA polymerase or ligase.

                    According to a convenient PCR method the nucleotide sequence encoding the  
30                   polypeptide of the invention and in particular the peptide addition thereof is used as a template and, e.g., doped or specific oligonucleotides are used as primers. In addition, cloning primers

localized outside the targetted region can be used. The resulting PCR product can either directly be cloned into an appropriate expression vector or gel purified and amplified in a second PCR reaction using the cloning primers and cloned into an appropriate expression vector.

In addition to the random mutagenesis methods described herein, it is occasionally useful to employ site specific mutagenesis techniques to modify one or more selected amino acids in the peptide addition, in particular to optimise the peptide addition with respect to the number of attachment groups.

Furthermore, random elongation mutagenesis as described by Matsuura et al, *op cit* can be used to construct a nucleotide sequence encoding a polypeptide having a C-terminal peptide addition. Construction of a nucleotide sequence encoding the polypeptide of the invention having an N-terminal peptide addition can be constructed in an analogous way.


Also, the methods disclosed in WO 97/04079, the contents of which are incorporated herein by reference, can be used for constructing a nucleotide sequence encoding the polypeptide of the invention.

The nucleotide sequence(s) or nucleotide sequence region(s) to be mutagenized is typically present on a suitable vector such as a plasmid or a bacteriophage, which as such is incubated with or otherwise exposed to the mutagenizing agent. The nucleotide sequence(s) to be mutagenized can also be present in a host cell either by being integrated into the genome of said cell or by being present on a vector harboured in the cell. Alternatively, the nucleotide sequence to be mutagenized is in isolated form. The nucleotide sequence is preferably a DNA sequence such as a cDNA, genomic DNA or synthetic DNA sequence.

Subsequent to the incubation with or exposure to the mutagenizing agent, the mutated nucleotide sequence, normally in amplified form, is expressed by culturing a suitable host cell carrying the nucleotide sequence under conditions allowing expression to take place. The host cell used for this purpose is one, which has been transformed with the mutated nucleotide sequence(s), optionally present on a vector, or one which carried the nucleotide sequence during the mutagenesis, or any kind of gene library.

#### DESIGN OF PEPTIDE ADDITION


One example of a useful guide for designing an N-terminal peptide addition containing N-glycosylation sites is characterized by the following formula:



wherein each of  $X_1$ ,  $X_3$  and  $X_4$  independently is absent or 1, 2, 3 or 4 amino acid residues of any type,  $X_2$  a single amino acid residue of any type except for proline,  $n$  any integer between 0 and 6, [T/S] a threonine or serine residue, preferably a threonine residue, and N and Pp has the meaning defined elsewhere herein. It has been found that sometimes the nature of the amino acid residue occupying position -1 to -4 relative to the N-residue of an N-glycosylation site may be important for the degree to which said N-glycosylation site is used. Accordingly,  $X_1$ ,  $X_3$ , and  $X_4$  may be chosen so as to obtain an increased utilization of the relevant site (as determined by a trial and error type of experiment). In a first step about 10 different muteins are made that has the above formula. For instance, the about 10 muteins are designed on the basis that each of  $X_1$ ,  $X_3$  and  $X_4$  independently is 1 or 2 alanine residues or is absent,  $Z$  any integer between 0 and 5, [T/S] threonine, and Alanine. Based on, e.g., *in vitro* bioactivity and half-life results obtained with these muteins (or any other relevant property), optimal number(s) of amino acids and glycosylation(s) can be determined and new muteins can be constructed based on this information. The process is repeated until an optimal glycosylated polypeptide is obtained.

Alternatively, random mutagenesis may be used for creating N-terminally extended polypeptides. For instance, a random mutagenized library is made on the basis of the above formula. Doped oligonucleotides are synthesized coding for one amino acid residue in position B (the amino acid residue being different from proline), each of  $X_1$ ,  $X_3$ , and  $X_4$  independently is 0, 1 or 2 amino acid residues of any type,  $n$  is 2 and T is threonine and used for constructing the random mutagenized library.

One example of a useful guide for designing an N-terminal peptide addition containing a PEGylation attachment group is characterized by the following formula using a lysine residue as an example of a PEGylation site. It will be understood that peptide additions with other attachment groups can be designed in an analogous way.



wherein each of  $Y^1$ ,  $Y^2$  and  $Y^3$  independently is 0, 1, 2, 3 or 4 amino acid residues of any type except lysine,  $n$  an integer between 0 and 6, K lysine, and Pp is as defined elsewhere herein.

In a first step about 10 different muteins are made that has the above formula. For instance, the about 10 muteins are designed on the basis that each of  $Y^1$ ,  $Y^2$  and  $Y^3$  independently is 1 or 2 alanine residues or is absent,  $n$  any integer between 0 and 5. The muteins are then PEGylated with 10 kDa PEG (e.g., using mPEG-SPA). Based on, e.g., *in vitro* bioactivity and

half-life results obtained with these muteins (or any other relevant property), optimal number(s) of amino acids and PEGylation sites can be determined and new muteins can be constructed based on this information. The process is repeated until an optimal PEGylated polypeptide is obtained.

Alternatively, random mutagenesis may be performed by making a random mutagenized library based on the above formula. Doped oligonucleotides are synthesized coding for one amino acid residue in position Y<sup>1</sup>, Y<sup>2</sup>, and/or Y<sup>3</sup> independently is 0, 1 or 2 amino acid residues of any type, and n is 2 and used for constructing the random mutagenized library.

#### Glucocerebrosidase (GCB) Activity Assay using PNP-glucopyranoside substrate

The enzymatic activity of recombinant GCB is measured using p-nitrophenyl-β-D-glucopyranoside (PNP-Glu) as a substrate. Hydrolysis of the PNP-Glu substrate generates p-nitrophenyl, which can be quantified by measuring absorption at 405 nm using a spectrophotometer, as previously described (Friedmann et al. (1999) *Blood* 93: 2807-2816). The assay is carried out under conditions which partially inhibit non-GCB glucosidase activities, such conditions being achieved by using a phosphate/citrate buffer pH 5.5, 0.25 % Triton X-100 and 0.25 % taurocholate.

The assay is run in a final volume of 200 μl, containing GCB Activity Assay Buffer and 4 mM PNP-Glu. The enzymatic hydrolysis is initiated by adding GCB and the reaction is allowed to proceed for 1 hour at 37°C before being stopped by adding 50 μl 1 M NaOH and measuring absorption at 405 nm. A reference standard curve of p-nitrophenyl, assayed in parallel, is used to quantify concentrations of GCB in samples to be tested.

#### In vitro uptake and stability of GCB polypeptide in macrophages

The murine monocyte/macrophage cells line, J774E (Mukhopadhyay and Stahl (1995) *Arch Biochem Biophys* 324(1): 78-84 and Diment et al. (1987) *J Leukoc Biol* 42(5): 485-90) is used to study the uptake and stability of GCB polypeptides. Cells are grown in alpha-MEM (supplemented with 10 % fetal calf serum, 1X Pen/Strep, and 60 μM 6-thioguanine), seeded (200,000 cells pr. well) in the above-mentioned media containing 10 μM conditol B epoxide, CBE (an irreversible GCB inhibitor) and incubated for 24 hr at 37°C.

Before starting the uptake assay, cells are washed in 0.5 ml HBSS (Hanks balanced salt solution). The uptake is done in a 200 μl volume, containing the appropriate concentration of GCB polypeptide (a dosis response curve is made with GCB concentrations in

the range of 25-400 mU/ml). As a control, yeast mannan (final concentration 1.4 mg/ml) is added to inhibit the uptake through the macrophage mannose receptor. The cells are incubated for 1 hr at 37°C and washed three times with 0.5 ml cold HBSS.

To measure the amount of GCB taken up by the J774E cells, cells are lysed in  
5 200 µl GCB Activity Assay Buffer with 4 mM PMP-Glu and incubated for 1 hr at 37°C. Then, the hydrolysis is stopped by addition of 50 µl 1M NaOH and OD405 is measured. The data are analysed by non-linear regression using GraphPad Prism 2.0 (GraphPad Software, San Diego, CA)

To study the stability of GCB polypeptides in J774E cells, CBE treated cells are  
10 incubated with 400 mU/ml GCB for 1 hr at 37°C. Then, cells are washed 3 times in HBSS to remove extracellular GCB and incubated in HBSS. A time-course study is done by lysing the cells after 30 min, 1 hr, 2 hr, 3hr, 4 hr, and 5 hr in 200 µl GCB Activity Assay Buffer with 4mM PNP-Glu and incubating the samples for 1 hr at 37°C before stopping the hydrolysis with 50 µl 1 M NaOH and measuring OD405. The data are analysed by non-linear regression using GraphPad  
15 Prism 2.0 (GraphPad Software, San Diego, CA).

#### Site-directed mutagenesis

Constructions of site-directed mutations were performed using PCR with  
oligonucleotides containing the desired amino acid exchanges or additions (e.g., to introduce glycosylation sites). The resulting PCR fragment was cloned into the GCB expression vector  
20 using appropriate restriction enzymes and subsequently DNA sequenced in order to confirm that the construct contained the desired exchanges.

#### MATERIALS

##### GCB Activity Assay Buffer:

120 mM phosphate/citrate buffer, pH=5.5, 1 mM EDTA, pH=8.0, 0.25 % Triton  
25 X-100, 0.25 % taurocholate, 4 mM β-mercaptoethanol.

##### pGC-12 vector

pVL1392 (Pharming, USA) with GCB wt cDNA sequence (SEQ ID NO 2)  
inserted between EcoRV and XbaI.

## TABLE 1

Sequence of primers used for cloning the wt GCB coding region and inserting signal peptides into the pGCBmat plasmid as described in Example 1.

- 5 SO49 (WT-sp-BglII): 5'-CGCAGATCTGATGGCTGGCAGCCTCACAGGATTGC-3'  
SO50 (WT-stop-EcoRI): 5'-CCGGAATTTCCATCACTGGCGACGCCACAGGTAGGTG-3'  
SO51 (WT-mature-SacI): 5'-ACGCGAGCTCGCCCCTGCATCCCTAAAAGCTTCGG-3'  
SO52 (SPegt-NheI/SacI-as): 5'-  
GCGTTGACGGCAGTCAGAGTTGACAGAAGGGCCAGCCAGCAAAGGATAGTCATG-3'  
10 SO53 (SPegt-NheI/SacI-s): 5'-  
CTAGCATGACTATCCTTTGCTGGCTGGCCCTTCTGTCAACTCTGACTGCCGTCAACGC  
AGCT-3'  
SO54 (SPegt-NheI/SacI-as): 5'-  
CCTGCTACTGCTCCCAGCAGCAGTGAAAGAGTCCAAAGTGGCAGCATG-3'  
15 SO55 (SPegt-NheI/SacI-s): 5'-  
CTAGCATGCTGCCACTTTGGACTCTTTCAGTCTGCTGGGAGCAGTAGCAGGAGCT-3'

*Cerezyme* was kindly provided by Dr. E. Beutler, Scripps Institute, CA, USA.

*J774E* was kindly provided by G. Grabowski, Cincinnati, Ohio, US

## EXAMPLE 1: PRODUCTION OF WT GCB

### Cloning and Expression in Insect Cells

20 A human fibroblast cDNA library was obtained from Clontech (Human fibroblast skin cDNA cloned in lambda-gt11, cat# HL1052b). Lambda DNA was prepared from the library by standard methods and used as a template in a PCR reaction with either SO49 and SO50 as primer (amplifies the GCB coding region with the human signal peptide from the second ATG)  
25 or SO50 and SO51 as primer (amplifies the mature part of the GCB coding region) (see Table 1 in the Materials section).

The PCR products were reamplified with the same primers and agarose gel purified. Subsequently the SO49/50 PCR product was digested with BglII and EcoRI and cloned into the pBlueBac 4.5 vector (Invitrogen, Carlsbad, CA, USA, Carlsbad, CA, USA)  
30 digested with BamHI and EcoRI. Sequencing confirmed that the insert is identical to the wtGCB

sequence as given in SEQ ID NO 2. The resulting plasmid was used for infection of insect cells with the GCB being partly secreted from the cells due to the human signal sequence as described in Martin et al., DNA 7, pp. 99-106, 1988. The SO50/51 PCR product was digested with SacI and EcoRI and cloned into the pBlueBac 4.5 vector (Invitrogen, Carlsbad, CA, USA) digested with the same enzymes resulting in the pGCBmat plasmid. Two different signal sequences were inserted upstream of the mature GCB codons in order to increase the secreted amount of enzyme. The baculovirus ecdysteroid UDPglucosyltransferase (egt) signal sequence (Murphy et al., Protein Expression and Purification 4, 349-357, 1993) was inserted by annealing SO52 and SO53 (Table 1) and the human pancreatic lipase signal sequence (Lowe et al., J. Biol. Chem. 264, 20042, 1989) was inserted by annealing SO54 and SO55 (Table 1) and cloning them into the NheI and SacI digested pGCBmat plasmid. Infection of *Spodoptera frugiperda* (Sf9) cells of the resulting plasmid was done according to the protocols from Invitrogen, Carlsbad, CA, USA.

#### Purification of GCB polypeptides produced in insect cells

Polypeptides with GCB activity were purified as described in US 5,236,838, with some modifications. Cells were removed from the culture medium by centrifugation (10 min at 4000 rpm in a Sorvall RC5C centrifuge) and the supernatant microfiltrated using a 0.22 µm filter prior to purification. DTT was added to 1 mM and the culture supernatant was ultrafiltrated to approximately 1/10 of the starting volume using a Vivaflow 200 system (Vivascience). The concentrated media was centrifuged to remove possible aggregates before application on a Toyopearl Butyl650C resin (TosoHaas) previously equilibrated in 50 mM sodium citrate, 20 % (v/v) ethylene glycol, 1 mM DTT, pH 5.0. This chromatographic step was performed at room temperature. The resin was washed with at least 3 column volumes of 50 mM sodium citrate, 20 % (v/v) ethylene glycol, 1 mM DTT, pH 5.0 (until the absorbance at 280 nm reaches baseline level) and GCB was eluted with a linear gradient from 0% to 100% 50 mM sodium citrate, 80% (v/v) ethylene glycol, 1 mM DTT, pH 5.0. Fractions were collected and assayed for GCB activity using the GCB Activity Assay. Usually, wt GCB starts to elute at approx. 70% (v/v) ethylene glycol.

The subsequent purification was done by either of the following two methods. #2 method results in GCB of a higher purity.

### Method #1

GCB enriched fractions from the first process step were pooled and diluted approx. 4 times with a buffer containing 50 mM sodium citrate, 5 mM DTT, pH 5.0 to reduce the ethylene glycol content to 20% (or lower). In the second HIC purification step the diluted and partially purified GCB was applied on a Toyopearl phenyl resin (TosoHaas) equilibrated in 50 mM sodium citrate, 1 mM DTT, pH 5.0 (Buffer A) before use. After application, the resin was washed with at least 3 column volumes of 50 mM sodium citrate, pH 5 (until the absorbance at 280 nm reaches baseline level) and GCB was then eluted with a linear ethanol gradient from 0% to 100% buffer B (50 mM sodium citrate, 50% (v/v) ethanol, 1 mM DTT, pH 5.0). Highly purified fractions of GCB (wildtype  $\geq 95\%$  pure), identified using the GCB Activity Assay, start to elute at approx. 40% ethanol. The purified GCB bulk product was dialyzed against 50 mM sodium citrate, 0.2 M mannitol, 0.09% tween80, pH 6.1 to retain the GCB activity upon subsequent storage at 4–8°C or at -80°C.

### Method #2

GCB enriched fractions eluted from the Toyopearl butyl650C resin were pooled and applied at 4°C on a SP sepharose resin (Amersham Pharmacia Biotech) previously equilibrated in 25 mM sodium citrate, 1 mM DTT, 10% ethylene glycol, pH 5.0. After application, the resin was washed with 25 mM sodium citrate, 1 mM DTT, 10% ethylene glycol, pH 5.0 (until absorption at 280 nm reached baseline level) and GCB was then eluted with a linear gradient from 0 to 100% 0.25 M sodium citrate, 1 mM DTT, 10% ethylene glycol, pH 5.0. GCB begins to elute around 0.15 M sodium citrate. Fractions containing GCB were pooled and applied at room temperature onto a Phenyl sepharose High Performance (Pharmacia Biotech) previously equilibrated in 25 mM sodium citrate 1 mM DTT, pH 5.0. After application, the resin was washed with 25 mM sodium citrate 1 mM DTT, pH 5.0 until absorption at 280 nm reached baseline level, and GCB was then eluted with a linear ethanol gradient from 0 to 100% 25 mM sodium citrate 1 mM DTT 50 % ethanol pH 5.0. GCB typically elutes around 35 % ethanol.

The purified GCB bulk product was dialyzed against either 50 mM sodium citrate, 1 mM DTT, pH 5.0 or 50 mM sodium citrate, 0.2 M mannitol, 1 mM DTT, pH 6.1 to retain the GCB activity upon subsequent storage. The purified GCB was concentrated and

sterilfiltered before storage at 4 – 8°C or at –80°C. Typically, GCB purified by this method is >95% pure.

## EXAMPLE 2

### Preparation of GCB with N-terminal peptide additions using a site-directed or random mutagenesis approach

Nucleotide sequences encoding the following N-terminal peptide additions were added to the nucleotide sequence shown in SEQ ID NO 2 encoding wtGCB: (A-4)+(N-3)+(I-2)+(T-1) (representing an extension to the N-terminal of the amino acid sequence shown in SEQ ID NO 1 with the amino acid residues ANIT), and (A-7)+(S-6)+(P-5)+(I-4)+(N-3)+(A-2)+(T-1) (ASPINAT).

A nucleotide sequence encoding the N-terminal peptide addition (A-4)+(N-3)+(I-2)+(T-1) was prepared by PCR using the following conditions:

#### PCR 1:

Template: 10 ng pBlueBac5 with wt GCB cDNA sequence

primer SO60: 5'-CAGCTGGCCATGGGTACCCGG-3' and

primer SO85:

5'-TGGGCATCAGGTGCCAACATTACAGCCCCGCCCTGCATCCCTAAAAGC-3'

BIO-X-ACT™ DNA polymerase (Bioline, London, U.K.)

1xOptiBuffer™ (Bioline, London, U.K.)

30 cycles of 96°C 30s, 55°C 30s, 72°C 1 min

#### PCR 2:

Template: 10 ng pBlueBac5 with wt GCB,

Baculo virus forward primer: 5'-TTTACTGTTTTTCGTAACAGTTTTG-3' and

PrimerSO86:

5'-GCAGGGGCGGGCTGTAATGTTGGCACCTGATGCCCACGACACTGCCTG-3'

BIO-X-ACT™ DNA polymerase (Bioline, London, U.K.)

1xOptiBuffer™ (Bioline, London, U.K.)

30 cycles of 96°C 30s, 55°C 30s, 72°C 1 min

### PCR 3:

3 µl of agarose gel purified PCR1 and PCR2 products (app. 10 ng)

Baculo virus forward primer: 5'-TTTACTGTTTTCGTAAACAGTTTGTG-3' and primer SO60.

BIO-X-ACT™ DNA polymerase (Bioline, London, U.K.)

5 1xOptiBuffer™ (Bioline, London, U.K.)

30 cycles of 96°C 30s, 55°C 30s, 72°C 1 min

PCR 3 was agarose gel purified and digested with NheI and NcoI and cloned into pBluebac4.5+wtGCB digested with NheI and NcoI.

After confirmation of the correct mutations by DNA sequencing the plasmid was  
10 transfected into insect cells using the Bac-N-Blue™ transfection kit from Invitrogen, Carlsbad, CA, USA. Expression of the muteins was tested by western blotting and by activity measurement of the muteins using the GCB Activity Assay.

Enzymatic activity of wtGCB (SEQ ID NO 1) expressed in the expression vector  
pVL1392 in insect cells (Sf9) using an analogous method to that described in Example 1 gave 13  
5 units/L, while the N-terminal peptide addition ASPINAT gave 28.5 units/L.

### Construction of libraries of GCB with N-terminal peptide addition

Using random mutagenesis two different libraries were constructed on the basis of  
GCB polypeptides with an N-terminal extension - library A with an N-terminal extension  
20 encoding the following amino acid sequence AXNXTXNXTXNXT, and library B with an N-terminal extension encoding ANXTNXTNXT.

Primers for library A were designed:

SO167: 5'-

GTGTCGTGGGCATCAGGTGCCNN(G/C)AA(C/T)(T/A/G)N(G/C)AC(A/T/C)(T/A/G)N(G/C)  
25 )AA(C/T)(T/A/G)N(G/C)AC(A/T/C)(T/A/G)N(G/C)AA(C/T)(T/A/G)N(G/C)AC(A/T/C)GCCC  
GCCCCTGCATCCCTAAAAGC

SO168 : 5'-GGCACCTGATGCCCACGACACTGCCTG

Primers for library B were designed using trinucleotides in the random positions.

X is a mixture of trinucleotide codons for all natural amino acid residues, except proline. The  
30 trinucleotide codons used were the same as described by Kayushin et al., Nucleic Acids  
Research, 24, 3748-3755, 1996.

SO165: 5'-

CGTGGGCATCAGGTGCCAAC(X)AC(A/T/C)AA(C/T)(X)AC(A/T/C)AA(C/T)(X)AC(A/T/C)  
)GCCCCGCCCTGCATCCCTAAAAGC

SO166 : 5'- GTTGGCACCTGATGCCCCACGACACTGCCTG

5 For both libraries:

SO60 and pBR10: 5'- TTT ACT GTT TTC GTA ACA GTT TTG

In all PCR reactions BIO-X-ACT™ DNA polymerase (Bioline, London, U.K.)  
and 1\*Optibuffer™ (Bioline, London, U.K.) were used. The PCR conditions were 30 cycles of  
10 94°C 30s, 55°C 1 min, and 72°C 1 min.

Templates and primers used for preparing a nucleotide sequence encoding the N-  
terminal extension by the above PCR were as follows:

PCR 1A:

Template: pGC12

Primers: SO60 + SO167

PCR 1B:

Template: pGC12

Primers: SO60 + SO165

20 PCR 2A:

Template: pGC12

Primers: SO168 + pBR10

PCR 2B:

Template: pGC12

25 Primers: SO166 + pBR10

PCR 3A:

Template: 1 µl of agarose gel purified PCR 1A and 2A products

Primers: SO60 + pBR10

### PCR 3B:

Template: 1 µl of agarose gel purified PCR 1B and 2B products

Primers: SO60 + pBR10

- PCR 3A and 3B were agarose gel purified and digested with NheI and NcoI and
- 5 ligated into pGC-12 digested with NheI and NcoI. The ligation mixture is transformed into competent *E. coli*. The diversity of the library was examined by DNA sequencing of different *E. coli* clones and gave rise to the following amino acid sequences:

### Library A:

- 1: AFNXTLNKTWN(F/L)T
- 10 2: TMNNTWNWTWNWT
- 3: -EXT wt
- 4: ALNSTGNLTVDGT
- 5: ASNSTFNLTENLT
- 6: TRNVTINCTUNST
- 15 7: -EXT wt
- 8: ALNWTYNGTKNVT
- 9: AANWTVNFTGNFT
- 10: -EXT wt
- 11: AXNXTVNSTUNVT
- 20 12: ANNFTFNGTLNLT
- 13: AGNWTANVTVNVT
- 14: AGNSTSNVTGNWT
- 15: AVNSTMNIHAIPP ( 1 deletion - nonsens)
- 16: AGNGTVNGTINGT
- 25 17: AVNSTGNXTGNWT
- 18: AGNGTUNGTSNLT
- 19: -EXT wt
- 20: AMNSTKNSTLNIT
- 21: AFNYTSKNST
- 30 22: -EXT wt
- 23: AVNATMNWTANGT

24: ASNSTNNGTLNAT  
 25: ARNKTKNFTINLT  
 26: APNITUNDTVNMT  
 27: AQNKTFNFTMNCT  
 5 28: ALNVTWNCTLNLT  
 29: ALNTTWTNLT

Library B:

1: ANTTNFTNET  
 2: ANWTNRTNCT  
 10 3: ANWTNFTNWT  
 4: PTGLIGNFT  
 5: ANWTNKTNFT  
 6: ANNTNLTNAT  
 7: ANYTNWTNFT  
 8: ANTTNQTNCT  
 9: - EXT wt  
 10: ANRTNWTNTT  
 11: PTATNHTNST  
 12: - EXT wt  
 13: ANWTNQTNQT  
 14: ANWTNWTNAT  
 15: ANFTNKTNMT  
 16: ANHTNETNAT  
 17: AN(C/W)TNFTNET  
 25 18: ANLDKLHKUH (insertion - nonsens)  
 19: ANCFTNQTNFT  
 20: ANWTNWTNEWT  
 21: ANCTNWTNCT  
 22: - EXT wt  
 30 23: - EXT wt  
 24: CHPYNWTNWT

- 25: ANETNYTNET
- 26: ANWTNWT
- 27: AKPYKSYKFY (insertion - nonsens)
- 28: ANITNKTNWT
- 5 29: ANWTNMTNIT
- 30: ANNTNRTNFT
- 31: ANWTNWTNWT
- 32: ANWRTNHTNKT
- 33: - EXT wt
- 10 34: ANQTNITNWT

Library B was transfected into insect cells using the Bac-N-Blue™ transfection kit from Invitrogen, Carlsbad, CA, USA. First, 96 plaques from Library B were picked and tested by activity measurement (GCB Activity Assay). Plaques were selected as follows: 3 with high activity, 3 with medium activity and 3 with low or no activity, and virus was purified for DNA sequencing resulting in the following amino acid sequences:

High activity:

- 1-1: Mixed sequence
- 1-2: ANFTNVATNQT
- 1-3: (A)(N)TTXLTN(K)T

Medium activity:

- 2-1: ANKTN(S/C)TNIT
- 2-2: Mixed sequence
- 2-3: ANWTNCTN(I)T

Low activity:

- 3-1: ANWTN(F/L)TNWT
- 3-2: CQLDURSTNET
- 3-3: No sequence

From both libraries 96 plaques were picked and tested by activity measurement (GCB Activity Assay). From each library 6 plaques with high activity were selected and virus

were purified for DNA sequencing. The amino acid sequence encoded by the different clones were:

Library A:

1: Mixed sequence

5 2: Mixed sequence

3: Mixed sequence

4: WT

5: ANNTNYTNWT

6: ANNTNYTNWT

10 Library B:

1: AANDTUNWTVNCT

2: ATNITLNYTANTT

3: WT

4: AANSTGNITINGT

15 5: AVNWTSNDTSNST

GCB polypeptides of the invention were tested for various properties, including GCB activity, stability in J774E cells and uptake in J774E cells. Unless otherwise stated the properties were tested by use of the methods described in the Methods section herein.

20 In the below table the GCB activity of various GCB polypeptides of the invention is listed together with the activity of the positives from Library A and B after plaque purification.

Table 2

25 Table 2: The plasmid column shows the number of the GCB polypeptide. The vector column shows the plasmid vector used for expression of the polypeptide. The mutation column shows the amino acid exchanges of the GCB polypeptide. N-terminal extensions are described as N-term followed by the amino acid residues that makes up the extension. The Activity column gives the units per liter of GCB activity measured by the GCB Activity Assay on the supernatant from Sf9 insect cells infected with one single plaque and grown in 3 ml of

media in a 6-well plate. Those labelled with P2 are activity measured of supernatant from virus infection cells grown in 15 ml T75 flasks.

Plasmid	Vector	Mutations	# Glycosylation sites introduced	Activity after Plaque Isolation (U/L)
pGC-1	PBlueBac4.5	Wt	0	6
pGC-6	pBlueBac4.5	N-termANIT	1	3
pGC-12	pVL1392	Wt	0	13
pGC-13	pVL1392	N-termASPINAT	1	29
pGC-36	pVL1392	N-term: ASPINATSPINAT	2	16
pGC-38	pVL1392	N-term: ASPINAT,K194N, K321N	3	16
pGC-40	pVL1392	N-term: ASPINAT,T132N, K293N, V295T	3	3.5
pGC-47	pVL1392	N-term: AGNGTVNGTINGT	3	30
pGC-48	pVL1392	N-term: ASNSTNNGTLNAT	3	36
pGC-56	pVL1392	N-term: ASPINATSPINAT, K194N, K321N	4	24
pGC-57	pVL1392	N-term: ASPINAT, T132N, K194N, K321N	4	20
pGC-58	pVL1392	N-term: ASPINAT, T132N, K194N	3	10
pGC-60	pVL1392	N-term:ANNTNYTNWT	3	P2: 14
pGC-61	pVL1392	N-term: ATNITLNYTANTT	3	P2: 38
pGC-62	pVL1392	N-term: AANSTGNITINGT	3	P2: 35
pGC-63	pVL1392	N-term: AVNWTSNDTSNST	3	P2: 66
pGC-68	pVL1392	AN N-term extension + R2T	1	37

**Table 3**

**Table 3:** Calculated  $V_{max}$  and  $K_m$  for uptake in the J774E macrophage cell line of the different GCB polypeptides.  $V_{max}$  and  $K_m$  was calculated from dosis response curve (See Fig. 1). The uptake of selected GCB polypeptides are shown in Figure 1.

As can be seen from table 3, an increase in  $V_{max}$  was observed for the N-terminally extended GCB polypeptides (pGC60, pGC61, and pGC62).

GCB polypeptide	Vmax	Km
Wildtype	0.57	87.7
Cerezyme	0.52	91.9
pGC36	0.60	70.6
pGC38	0.48	44.0
pGC56	0.39	32.2
pGC60	0.57	79.1
pGC61	0.74	100.5
pGC62	0.86	110.8
pGC63	0.51	83.1

### EXAMPLE 3: GLYCOSYLATION OF GCB POLYPEPTIDES OF THE INVENTION EXPRESSED IN INSECT CELLS

MALDI-TOF mass spectrometry was used to investigate the amount of carbohydrate attached to GCB polypeptides expressed in Sf9 cells.

The 6 GCB polypeptide variants investigated all contained additional potential N-glycosylation sites compared to wtGCB.

WtGCB contains 5 potential N-glycosylation sites of which only 4 are used.

The 6 GCB polypeptide variants were:

- GC-36: ASPINATSPINAT-GCB,
- GC-38: ASPINAT-GCB(K194N,K321N),
- GC-60: ANNTNYTNWT-GCB,
- GC-61: ATNITLNYTANTT-GCB,
- GC-62: AANSTGNITINGT-GCB, and
- GC-63: AVNWTSNDTSNST-GCB.

#### WtGCB:

The theoretical peptide mass of wtGCB is 55 591 Da. WtGCB has 5 potential N-glycosylation sites of which only 4 are used. As the two most common N-glycan structures on recombinant proteins expressed in Sf9 cells are Man<sub>3</sub>GlcNAc<sub>2</sub>Fuc and Man<sub>3</sub>GlcNAc<sub>2</sub> having masses of 1038.38 Da and 892.31 Da, respectively, the expected mass of wtGCB carrying 4 N-glycans is between 59 159 Da and 59 743 Da.

MALDI-TOF mass spectrometry of wtGCB shows the broad peak typical of glycoproteins with a peak mass of 59.3 kDa in accordance with the expected mass of wtGCB carrying 4 N-glycans.

GC-36 (ASPINATSPINAT-GCB):

The theoretical peptide mass of GC-36 is 56 829 Da. The N-terminal extension contains two additional potential glycosylation sites at N5 and N11 compared to wtGCB. Assuming that the wtGCB part of the variant is glycosylated like wtGCB, the variant has 6 potential N-glycosylation sites.

As the two most common N-glycan structures on recombinant proteins expressed in Sf9 cells are Man<sub>3</sub>GlcNAc<sub>2</sub>Fuc and Man<sub>3</sub>GlcNAc<sub>2</sub> having masses of 1038.38 Da and 892.31 Da, respectively, the expected mass of GC-36 carrying 4 N-glycans is between 60 397 Da and 60 981 Da, the expected mass of GC-36 carrying 5 N-glycans is between 61 289 Da and 62 019 Da, and the expected mass of GC-36 carrying 6 N-glycans is between 62 181 Da and 63 057 Da.

MALDI-TOF mass spectrometry of GC-36 shows a rather broad peak with a peak mass between 61.5 kDa and 62.9 kDa in accordance with the expected mass of GC-36 carrying either 5 or 6 N-glycans.

N-terminal amino acid sequence analysis of GC-36 showed that N5 is completely glycosylated while N11 is partially glycosylated in complete agreement with the result obtained using mass spectrometry.

GC-38 (ASPINAT-GCB(K194N,K321N)):

The theoretical peptide mass of GC-38 is 56 217 Da. The N-terminal extension contains one additional potential glycosylation sites at N5 compared to wtGCB. In addition, the substitutions of Lys194 and Lys321 with Asn-residues introduce two additional potential N-glycosylation sites. Assuming that the wtGCB part of the variant is glycosylated like wtGCB, the variant has 7 potential N-glycosylation sites.

Based on the same considerations as those used for GC-36, the expected mass of GC-38 carrying 4 N-glycans is between 59 785 Da and 60 369 Da, the expected mass of GC-38 carrying 5 N-glycans is between 60 677 Da and 61 407 Da, the expected mass of GC-38 carrying 6 N-glycans is between 61 569 Da and 62 445 Da, and the expected mass of GC-38 carrying 7 N-glycans is between 62 461 Da and 63 483 Da.

MALDI-TOF mass spectrometry of GC-38 shows a major peak with a peak mass of 63.1 kDa in accordance with the expected mass of GC-38 carrying 7 N-glycans. In addition, a minor peak with a peak mass of 62.3 kDa is seen which corresponds to GC-38 carrying 6 N-glycans.

5 N-terminal amino acid sequence analysis of GC-38 showed that N5 is completely glycosylated.

GC-60 (ANNTNYTNWT-GCB):

The theoretical peptide mass of GC-60 is 56 770 Da. The N-terminal extension contains three additional potential glycosylation sites at N2, N5 and N8 compared to wtGCB.

10 Assuming that the wtGCB part of the variant is glycosylated like wtGCB, the variant has 7 potential N-glycosylation sites.

Based on the same considerations as those used for GC-36 the expected mass of GC-60 carrying 4 N-glycans is between 60 338 Da and 60 922 Da, the expected mass of GC-60 carrying 5 N-glycans is between 61 230 Da and 61 960 Da, the expected mass of GC-60 carrying 6 N-glycans is between 62 122 Da and 62 998 Da, and the expected mass of GC-60 carrying 7 N-glycans is between 63 014 Da and 64 036 Da.

MALDI-TOF mass spectrometry of GC-60 shows two broad peaks with peak masses of 61.9 kDa and 62.8 kDa in accordance with the expected mass of GC-60 carrying either 5 or 6 N-glycans.

20 N-terminal amino acid sequence analysis of GC-60 showed that N2 is mainly glycosylated, N5 is completely glycosylated while N8 is only seldom glycosylated in acceptable agreement with the result obtained using mass spectrometry.

GC-61 (ATNITLNYTANTT-GCB):

25 The theoretical peptide mass of GC-61 is 56 970 Da. The N-terminal extension contains three additional potential glycosylation sites at N3, N7 and N11 compared to wtGCB. Assuming that the wtGCB part of the variant is glycosylated like wtGCB, the variant has 7 potential N-glycosylation sites.

30 Based on the same considerations as used for GC-36, the expected mass of GC-61 carrying 4 N-glycans is between 60 538 Da and 61 122 Da, the expected mass of GC-61 carrying 5 N-glycans is between 61 430 Da and 62 160 Da, the expected mass of GC-61 carrying 6 N-

glycans is between 62 322 Da and 63 198 Da, and the expected mass of GC-61 carrying 7 N-glycans is between 63 214 Da and 64 236 Da.

MALDI-TOF mass spectrometry of GC-61 shows a very broad peak with peak mass between 61.5 kDa and 63.0 kDa in accordance with the expected mass of GC-61 carrying either 5 or 6 N-glycans.

N-terminal amino acid sequence analysis of GC-61 showed that N3 is completely glycosylated while N7 and N11 are partially glycosylated in acceptable agreement with the result obtained using mass spectrometry.

#### GC-62 (AANSTGNITINGT-GCB):

The theoretical peptide mass of GC-62 is 56 806 Da. The N-terminal extension contains three additional potential glycosylation sites at N3, N7 and N11 compared to wtGCB. Assuming that the wtGCB part of the variant is glycosylated like wtGCB, the variant has 7 potential N-glycosylation sites.

Based on the same considerations as those used for GC-36, the expected mass of GC-62 carrying 4 N-glycans is between 60 374 Da and 60 958 Da, the expected mass of GC-62 carrying 5 N-glycans is between 61 266 Da and 61 996 Da, the expected mass of GC-62 carrying 6 N-glycans is between 62 158 Da and 63 034 Da, and the expected mass of GC-62 carrying 7 N-glycans is between 63 050 Da and 64 072 Da.

MALDI-TOF mass spectrometry of GC-62 shows two broad peaks with peak masses of 61.6 kDa and 62.7 kDa in accordance with the expected mass of GC-62 carrying either 5 or 6 N-glycans.

N-terminal amino acid sequence analysis of GC-62 showed that N3 is completely glycosylated while N7 and N11 are partially glycosylated in acceptable agreement with the result obtained using mass spectrometry.

#### GC-63 (AVNWTSNDTSNST-GCB):

The theoretical peptide mass of GC-63 is 56 969 Da. The N-terminal extension contains three additional potential glycosylation sites at N3, N7 and N11 compared to wtGCB. Assuming that the wtGCB part of the variant is glycosylated like wtGCB, the variant has 7 potential N-glycosylation sites.

Based on the same considerations as those used for GC-36, the expected mass of GC-63 carrying 4 N-glycans is between 60 537 Da and 61 121 Da, the expected mass of GC-63

carrying 5 N-glycans is between 61 429 Da and 62 159 Da, the expected mass of GC-63 carrying 6 N-glycans is between 62 321 Da and 63 197 Da, and the expected mass of GC-63 carrying 7 N-glycans is between 63 213 Da and 64 235 Da.

MALDI-TOF mass spectrometry of GC-63 shows a major peak with a peak mass of 61.9 kDa in accordance with the expected mass of GC-63 carrying 5 N-glycans. In addition, a minor peak with a peak mass of 62.9 kDa is seen which corresponds to GC-63 carrying 6 N-glycans.

N-terminal amino acid sequence analysis of GC-63 showed that N3 and N7 are partially glycosylated. It was not possible to evaluate the glycosylation status of N11.

Furthermore, insect cell expressed N-terminally extended glycosylated polypeptide (GC-6 and GC-13) was subjected to N-terminal amino acid sequence analysis (using Procize from PE Biosystems, Foster City, CA). The sequencing cycle was blank for the Asn residue in both ANIT and ASPINAT N-terminal peptide additions, demonstrating that the introduced glycosylation site is glycosylated.

When subjecting GC-13 to mass spectrophotometry using the MALDI-TOF techniques on the Voyager DERP instrument (from PE-Biosystems, Foster City, CA) the following results were obtained:

The wildtype and ASPINAT-extended wildtype expressed in insect cells gave average masses very close to the calculated mass of 59,727 Da and 61,421 Da, respectively, assuming that four glycosylation sites were occupied by the carbohydrates FucGlcNAc<sub>2</sub>Man<sub>3</sub>.

#### EXAMPLE 4: CONSTRUCTION OF PLASMIDS FOR EXPRESSION OF FSH

A gene encoding the human FSH-alpha subunit was constructed by assembly of synthetic oligonucleotides by PCR using methods similar to the ones described in Stemmer et al. (1995) *Gene* 164, pp. 49-53. The native FSH-alpha signal sequence was maintained in order to allow secretion of the gene product. The codon usage of the gene was optimised for high expression in mammalian cells. Furthermore, in order to achieve high gene expression, an intron (from pCI-Neo (Promega)) was included in the 5' untranslated region of the gene. The synthetic gene was subcloned behind the CMV promoter in pcDNA3.1/Hygro (Invitrogen). The sequence of the resulting plasmid, termed pBvdH977, is given in SEQ ID NO:3 (FSH-alpha-coding sequence at position 1225 to 1570). Similarly, a synthetic gene encoding the wildtype human FSH-beta subunit was constructed. Also in this construct, the native signal sequence was

maintained (except for a Lys to Glu mutation at position 2) in order to allow secretion, and the codon usage was optimised for high expression and an intron was included in the recipient vector (pcDNA3.1/Zeo (Invitrogen)). The sequence of the resulting FSH-beta -containing plasmid, termed pBvdH1022, is given in SEQ ID NO:4 (FSH-beta-coding sequence at position 1231 to 1617). A plasmid containing both the FSH-alpha and the FSH-beta encoding synthetic genes was generated by subcloning the FSH-alpha containing *NruI*-*PvuII* fragment from pBvdH977 into pBvdH1022 linearized with *NruI*. The resulting plasmid, in which the FSH-alpha and FSH-beta-expression cassettes are in direct orientation, was termed pBvdH1100.

#### Expression of FSH in CHO cells

FSH was expressed in Chinese Hamster Ovary (CHO) K1 cells, obtained from the American Type Culture Collection (ATCC, CCL-61).

For transient expression of FSH, cells were grown to 95% confluency in serum-containing media (MEM $\alpha$  with ribonucleotides and deoxyribonucleotides (Life Technologies Cat # 32571-028) containing 1:10 FBS (BioWhittaker Cat # 02-701F) and 1:100 penicillin and streptomycin (BioWhittaker Cat # 17-602E), or Dulbecco's MEM/Nut.-mix F-12 (Ham) L-glutamine, 15 mM Hepes, pyridoxine-HCl (Life Technologies Cat # 31330-038) with the same additives. FSH-encoding plasmids were transfected into the cells using Lipofectamine 2000 (Life Technologies) according to the manufacturer's specifications. 24-48 hrs after transfection, culture media were collected, centrifuged and filtered through 0.22 micrometer filters to remove cells.

Stable clones expressing FSH were generated by transfection of CHO K1 cells with FSH-encoding plasmids followed by incubation of the cells in selective media (for instance one of the above media containing 0.5 mg/ml zeocin for cells transfected with plasmid pBvdH1100). Stably transfected cells were isolated and sub-cloned by limited dilution. Clones that produced high levels of FSH were identified by ELISA.

More specifically, the concentration of FSH in samples was quantified by use of a commercial immunoassay (DRG FSH EIA, DRG Instruments GmbH, Marburg, Germany). DRG FSH EIA is a solid phase immunosorbent assay (ELISA) based on the sandwich principle. The microtiter wells are coated with a monoclonal antibody directed towards a unique antigenic site on the FSH- $\beta$  subunit. An aliquot of FSH-containing sample (diluted in H<sub>2</sub>O with 0.1% BSA) and an anti-FSH antiserum conjugated with horseradish peroxidase are added to the coated wells.

After incubation, unbound conjugate is washed off with water. The amount of bound peroxidase is proportional to the concentration of FSH in the sample. The intensity of colour developed upon addition of substrate solution is proportional to the concentration of FSH in the sample.

#### Large-scale production of FSH in CHO cells

The cell line CHO K1 1100-5, stably expressing human FSH, was passed 1:10 from a confluent culture and propagated as adherent cells in serum-containing medium Dulbecco's MEM/Nut.-mix F-12 (Ham) L-glutamine, 15 mM Hepes, pyridoxine-HCl (Life Technologies Cat # 31330-038), 1:10 FBS (BioWhittaker Cat # 02-701F), 1:100 penicillin and streptomycin (BioWhittaker Cat # 17-602E) until confluence in a 10 layer cell factory (NUNC #165250). The media was then changed to serum-free media: Dulbecco's MEM/Nut.-mix F-12 (Ham) L-glutamine, pyridoxine-HCl (Life Technologies Cat # 21041-025) with the addition of 1:500 ITS-A (Gibco/BRL # 51300-044), 1:500 EX-CYTE VLE (Serological Proteins Inc. # 81-129) and 1:100 penicillin and streptomycin (BioWhittaker Cat # 17-602E). Subsequently, every 24 h, culture media were collected and replaced with 1 fresh liter of the same serum-free media. The collected media was filtered through 0.22 µm filters to remove cells. Growth in cell factories was continued with daily harvests and replacements of the culture media until FSH yields dropped below one-fourth of the initial expression level (typically after 10-15 days).

#### EXAMPLE 5: PURIFICATION OF FSH WILDTYPE AND VARIANTS

Three chromatographic steps were employed to obtain highly purified FSH. First an anion exchanger step, then hydrophobic interaction chromatography (HIC) and finally an immunoaffinity step using an FSH-β specific monoclonal antibody.

Culture supernatants were prepared as described in Example 4. Filtered culture supernatants were concentrated 10 to 20 times by ultrafiltration (10 kD cut-off membrane), pH was adjusted to 8.0 and conductivity to 10 - 15 mS/cm, before application on a DEAE Sepharose (Pharmacia) anion exchanger column, which had been equilibrated in ammonium acetate buffer (0.16 M, pH 8.0). Semipurified FSH was recovered both in the unbound flow-through fraction as well as in the wash fraction using 0.16 M ammonium acetate, pH 8.0. The flow through and wash fractions were pooled and ammonium sulfate was added from a stock solution (4.5 M) to obtain a final concentration of 1.5 M (NH<sub>4</sub>)<sub>2</sub>SO<sub>4</sub>. The pH was adjusted to 7.0.

The partially purified FSH was subsequently applied on a 25 ml butyl Sepharose (Pharmacia) HIC column. After application, the column was washed with at least 3 column volumes of 1.5 M  $(\text{NH}_4)_2\text{SO}_4$ , 20 mM ammonium acetate, pH 7 (until the absorbance at 280 nm reached baseline level) and FSH was eluted with 4 column volumes of buffer B (20 mM ammonium acetate, pH 7). FSH enriched fractions from the HIC step were pooled, concentrated and diafiltrated using Vivaspin 20 modules, 10 kD cut-off membrane (Vivascience), to a 50 mM sodium phosphate, 150 mM NaCl, pH 7.2.

For the third chromatographic step, an anti-FSH- $\beta$  monoclonal antibody (RDI-FSH909, Research Diagnostics) was immobilized to CNBr-activated Sepharose (Pharmacia) using a standard procedure from the supplier. Approximately 1 mg antibody was coupled per ml resin. The immunoaffinity resin was packed in plastic columns and equilibrated with 50 mM sodium phosphate, 150 mM NaCl, pH 7.2 before application.

The buffer exchanged eluate from the butyl HIC step was applied on the antibody column by use of gravity flow. This was followed by several washing steps in 50 mM sodium phosphate solutions (0.5 M NaCl and 1 M NaCl, both pH 7.2). Elution was performed using either 1 M  $\text{NH}_3$  or 0.6 M  $\text{NH}_3$ , 40% (v/v) isopropanol and the eluate was immediately neutralized with 1 M acetic acid to pH 6-8.

The purified FSH bulk product was concentrated and diafiltrated using Vivaspin 20 modules, 10 kD cut-off membrane (Vivascience), to a 50 mM sodium phosphate, 150 mM NaCl, pH 7.2. For subsequent storage, BSA was added to 0.1% (w/v) and the purified FSH was microfiltrated using a 0.22  $\mu\text{m}$  filter prior to storage at  $-80^\circ\text{C}$ .

SDS-PAGE, run under non-dissociating conditions (without boiling), showed wildtype FSH migrating as an apparant  $42\pm3$  kDa band, slightly diffuse due to heterogeneity in the attached carbohydrates. The purity was about 80-90%. N-terminal sequencing showed that the  $\alpha$ -chain had the expected N-terminal sequence starting with residue 1 (SEQ ID NO:5) and the  $\beta$ -chain starting with residue 3 (SEQ ID NO:6). These N-terminal sequences have been found previously for recombinant FSH produced in CHO cells (Olijve, W. et al. (1996) *Mol. Hum. Reprod.* 2, 371-382).

## EXAMPLE 6 FSH IN VITRO ACTIVITY ASSAY

### 6.1 FSH assay Outline

It has previously been published that activation of the FSH receptor by FSH leads to an increase in the intracellular concentration of cAMP. Consequently, transcription is activated at promoters containing multiple copies of the cAMP response element (CRE). It is thus possible to measure FSH activity by use of a CRE luciferase reporter gene introduced into CHO cells expressing the FSH receptor.

### 6.2 Construction of a CHO FSH-R / CRE-luc cell line

Stable clones expressing the human FSH receptor were produced by transfection of CHO K1 cells with a plasmid containing the receptor cDNA inserted into pcDNA3 (Invitrogen) followed by selection in media containing 600 microg/ml G418. Using a commercial cAMP-SPA RIA (Amersham), clones were screened for the ability to respond to FSH stimulation. On the basis of these results, an FSH receptor-expressing CHO clone was selected for further transfection with a CRE-luc reporter gene. A plasmid containing the reporter gene with 6 CRE elements in front of the Firefly luciferase gene was co-transfected with a plasmid conferring Hygromycin B resistance. Stable clones were selected in the presence of 600 microg/ml G418 and 400 microg/ml Hygromycin B. A clone yielding a robust luciferase signal upon stimulation with FSH ( $EC_{50} \sim 0.01$  IU/ml) was obtained. This CHO FSH-R / CRE-luc cell line was used to measure the activity of samples containing FSH.

### 6.3 FSH luciferase assay

To perform activity assays, CHO FSH-R / CRE-luc cells were seeded in white 96 well culture plates at a density of about 15,000 cells/well. The cells were in 100  $\mu$ l DMEM/F-12 (without phenol red) with 1.25% FBS. After incubation overnight (at 37°C, 5% CO<sub>2</sub>), 25  $\mu$ l of sample or standard diluted in DMEM/F-12 (without phenol red) with 10% FBS was added to each well. The plates were further incubated for 3 hrs, followed by addition of 125  $\mu$ l LucLite substrate (Packard Bioscience). Subsequently, plates were sealed and luminescence was measured on a TopCount luminometer (Packard) in SPC (single photon counting) mode.

EXAMPLE 7: CONSTRUCTION AND ANALYSIS OF A VARIANT FORM OF FSH CONTAINING TWO N-LINKED GLYCOSYLATIONS AT THE N-TERMINUS OF THE ALPHA SUBUNIT

A construct encoding a modified form of FSH-alpha, having two additional sites for N-linked glycosylation at its N-terminus was generated by site-directed mutagenesis using standard DNA techniques known in the art. A DNA fragment encoding the sequence Ala-Asn-Ile-Thr-Val-Asn-Ile-Thr-Val was inserted immediately upstream of the mature FSH-alpha sequence in pBvdH977. The sequence of the resulting plasmid, termed pBvdH1163, is given in SEQ ID NO:7 (modified FSH-alpha-encoding sequence at position 1225 to 1599). A plasmid encoding both subunits was constructed by subcloning the FSH-containing *NruI*-*PvuII* fragment from pBvdH1163 into pBvdH1022 (Example 4), which had been linearized with *PvuII*. The resulting plasmid was termed pBvdH1208.

For expression of the variant form of FSH containing two N-linked glycosylations at the N-terminus of the alpha subunit (termed FSH1208), CHO K1 cells were transfected with pBvdH1208 or co-transfected with a combination of pBvdH1163, encoding the modified alpha subunit and pBvdH1022, encoding the wildtype beta subunit. Transient expressions, isolation of stable expression clones, and large-scale production of FSH1208 were performed as described for wildtype FSH in Example 4.

The FSH content of samples was analysed by Western Blotting: Proteins were separated by SDS-PAGE and a standard Western blot was performed using rabbit anti human FSH (AHP519, Serotec) or mouse anti human FSH-alpha (MCA338, Serotec) as primary antibody, and an ImmunoPure Ultra Sensitive ABC Peroxidase Staining Kit (Pierce) for detection. Western blotting showed that FSH1208 had a larger molecular mass than wildtype FSH, indicating that the introduction of acceptor sites for N-linked glycosylation at the N-terminus of the alpha subunit indeed lead to hyperglycosylation of FSH. For analysis of pI, samples were separated on pH 3-7 IEF gels (NOVEX). After electrophoresis, proteins were blotted onto Immobilon-P (Millipore) membranes and a Western blot was performed as described above, using the same antibodies and detection kit. Isoelectric focusing demonstrated that the FSH forms in the FSH1208 samples were found in a lower pI range than wildtype FSH. Thus, the pH interval for FSH1208 isoforms was about 3.0-4.5 versus about 4.0-5.2 for wildtype FSH. This indicated that FSH1208 molecules are on average more negatively charged than the wild type, which is attributed to the presence of additional sialic acid residues.

FSH1208 was purified and characterized as described in Example 5. SDS-PAGE, run under non-dissociating conditions (without boiling), showed FSH1208 migrating as an apparent 55±5 kDa band, slightly diffuse due to heterogeneity in the attached carbohydrates. The purity was about 80-90%. N-terminal sequencing showed that while the β-chain had the same N-terminal sequence as wildtype FSH, the sequence of α-chain was in agreement with this subunit carrying the expected N-terminal extension ANITVNITV, in which both asparagines residues are glycosylated.

The specific activity of FSH1208 was determined by measurement of the *in vitro* bioactivity (FSH luciferase assay, Example 6) and the FSH content of the samples by ELISA.

The specific activity of FSH1208 was found to be about one-third of that of the wildtype reference.

A pharmacokinetic study performed as follows:

Immature 26-27 days old female Sprague-Dawley rats were injected i.v. with 3-4 microg FSH, produced, purified and analyzed as described above. Subsequently, blood samples were taken at various time-points after injection. FSH concentrations in serum samples were determined by ELISA, as described above.

*In vivo* bioactivity of wildtype recombinant FSH and variant forms may be evaluated by the ovarian weight augmentation assay (Steelman and Pohley (1953) *Endocrinology* **53**, 604-616). Furthermore, the ability of FSH and variant forms to stimulate maturation of follicles in laboratory animals may be detected with e.g., ultrasound equipment. The experiment showed that 24 hours after injection of equal amounts of wildtype FSH and FSH1208, the sera of FSH1208-treated animals contained more than 10 fold more remaining immunoreactive material than the sera from animals treated with wildtype FSH.

#### EXAMPLE 8: CONSTRUCTION AND ANALYSIS OF OTHER FSH VARIANTS CONTAINING ADDITIONAL GLYCOSYLATION SITES

Plasmids encoding variant forms of FSH-alpha and FSH-beta containing additional sites for N-linked glycosylation were generated by site-directed mutagenesis using standard DNA techniques known in the art. The following amino acid substitutions and/or insertions were generated:

FSH1147: Amino acid Tyr58 of mature FSH-beta altered to Asn

FSH1349: N-terminus of mature FSH-alpha altered from APD QDC... to: APNDTVNFT QDC

...

FSH1354: N-terminus of mature FSH-beta altered from NS CEL ... to: NSNITVNITV CEL ...

Plasmids encoding the variant forms were transiently expressed in CHO K1 cells  
5 as described in Example 4. Plasmids encoding FSH-alpha variants were co-transfected with a  
plasmid encoding wild-type FSH-beta and vice versa.

Western and isoelectric focusing were performed on culture media samples as  
described above. The variant forms had higher molecular weights than the wild-type, indicating  
that the additional acceptor sites for N-linked glycosylation had indeed been glycosylated.

10 Furthermore, isoelectric focusing showed that the different isoforms of the three FSH variants  
were spread over a lower pI range than the wildtype. This strongly suggests that the variant  
forms had a higher sialic acid content than the wildtype.

*In vitro* FSH activities of the resulting media samples were analysed as described  
in Example 6.3. All three variant forms were able to stimulate the CHO FSH-R / CRE-luc cells,  
15 indicating that these variant FSH forms have retained significant FSH activity.

While the foregoing invention has been described in some detail for purposes of  
clarity and understanding, it will be clear to one skilled in the art from a reading of this  
disclosure that various changes in form and detail can be made without departing from the true  
scope of the invention. For example, all the techniques, methods, compositions, apparatus and  
20 systems described above may be used in various combinations. All publications, patents, patent  
applications, or other documents cited in this application are incorporated by reference in their  
entirety for all purposes to the same extent as if each individual publication, patent, patent  
application, or other document were individually indicated to be incorporated by reference for all  
purposes.

25

# SEQUENCE LISTING

<110> MAXYGEN APS

<120> N-TERMINALLY EXTENDED POLYPEPTIDES

<130> 0217WO210

<170> PatentIn Ver. 2.1

<210> 1

<211> 497

<212> PRT

<213> Homo sapiens

<220>

<221> MOD\_RES

<222> (495)

<223> R or H

<400> 1

Ala Arg Pro Cys Ile Pro Lys Ser Phe Gly Tyr Ser Ser Val Val Cys  
1 5 10 15

Val Cys Asn Ala Thr Tyr Cys Asp Ser Phe Asp Pro Pro Thr Phe Pro  
20 25 30

Ala Leu Gly Thr Phe Ser Arg Tyr Glu Ser Thr Arg Ser Gly Arg Arg  
35 40 45

Met Glu Leu Ser Met Gly Pro Ile Gln Ala Asn His Thr Gly Thr Gly  
50 55 60

Leu Leu Leu Thr Leu Gln Pro Glu Gln Lys Phe Gln Lys Val Lys Gly  
65 70 75 80

Phe Gly Gly Ala Met Thr Asp Ala Ala Ala Leu Asn Ile Leu Ala Leu  
85 90 95

Ser Pro Pro Ala Gln Asn Leu Leu Leu Lys Ser Tyr Phe Ser Glu Glu  
100 105 110

Gly Ile Gly Tyr Asn Ile Ile Arg Val Pro Met Ala Ser Cys Asp Phe  
115 120 125

Ser Ile Arg Thr Tyr Thr Tyr Ala Asp Thr Pro Asp Asp Phe Gln Leu  
130 135 140

His	Asn	Phe	Ser	Leu	Pro	Glu	Glu	Asp	Thr	Lys	Leu	Lys	Ile	Pro	Leu	145	150	155	160
Ile	His	Arg	Ala	Leu	Gln	Leu	Ala	Gln	Arg	Pro	Val	Ser	Leu	Leu	Ala	165	170	175	
Ser	Pro	Trp	Thr	Ser	Pro	Thr	Trp	Leu	Lys	Thr	Asn	Gly	Ala	Val	Asn	180	185	190	
Gly	Lys	Gly	Ser	Leu	Lys	Gly	Gln	Pro	Gly	Asp	Ile	Tyr	His	Gln	Thr	195	200	205	
Trp	Ala	Arg	Tyr	Phe	Val	Lys	Phe	Leu	Asp	Ala	Tyr	Ala	Glu	His	Lys	210	215	220	
Leu	Gln	Phe	Trp	Ala	Val	Thr	Ala	Glu	Asn	Glu	Pro	Ser	Ala	Gly	Leu	225	230	235	240
Leu	Ser	Gly	Tyr	Pro	Phe	Gln	Cys	Leu	Gly	Phe	Thr	Pro	Glu	His	Gln	245	250	255	
Arg	Asp	Phe	Ile	Ala	Arg	Asp	Leu	Gly	Pro	Thr	Leu	Ala	Asn	Ser	Thr	260	265	270	
His	His	Asn	Val	Arg	Leu	Leu	Met	Leu	Asp	Asp	Gln	Arg	Leu	Leu	Leu	275	280	285	
Pro	His	Trp	Ala	Lys	Val	Val	Leu	Thr	Asp	Pro	Glu	Ala	Ala	Lys	Tyr	290	295	300	
Val	His	Gly	Ile	Ala	Val	His	Trp	Tyr	Leu	Asp	Phe	Leu	Ala	Pro	Ala	305	310	315	320
Lys	Ala	Thr	Leu	Gly	Glu	Thr	His	Arg	Leu	Phe	Pro	Asn	Thr	Met	Leu	325	330	335	
Phe	Ala	Ser	Glu	Ala	Cys	Val	Gly	Ser	Lys	Phe	Trp	Glu	Gln	Ser	Val	340	345	350	
Arg	Leu	Gly	Ser	Trp	Asp	Arg	Gly	Met	Gln	Tyr	Ser	His	Ser	Ile	Ile	355	360	365	
Thr	Asn	Leu	Leu	Tyr	His	Val	Val	Gly	Trp	Thr	Asp	Trp	Asn	Leu	Ala	370	375	380	
Leu	Asn	Pro	Glu	Gly	Gly	Pro	Asn	Trp	Val	Arg	Asn	Phe	Val	Asp	Ser				



```

caccgagcac tgcagttggc ccagcgtccc gtttcactcc ttgccagccc
ctggacatca 600
cccacttggc tcaagaccaa tggagcgggtg aatgggaagg ggtcactcaa
gggacagccc 660
ggagacatct accaccagac ctggggccaga tactttgtga agttcctgga
tgcctatgct 720
gagcacaagt tacagttctg ggcagtgaca gctgaaaatg agccttctgc
tgggctggtg 780
agtggatacc ccttccagtg cctgggcttc acccctgaac atcagcgaga
cttaattgcc 840
cgtgacctag gtcctaccct cgccaacagt actcaccaca atgtccgcct
actcatgctg 900
gatgaccaac gcttgctgct gccccactgg gcaaagggtgg tgctgacaga
cccagaagca 960
gctaaatatg ttcattggcat tgctgtacat tggtagctgg actttctggc
tccagccaaa 1020
gccaccctag gggagacaca ccgcctgttc cccaacacca tgctctttgc
ctcagaggcc 1080
tgtgtgggct ccaagttctg ggagcagagt gtgaggctag gctcctggga
tcgagggatg 1140
cagtacagcc acagcatcat cacgaacctc ctgtaccatg tggtcggctg
gaccgactgg 1200
aaccttgccc tgaaccccga aggaggacce aattgggtgc gtaactttgt
cgacagtccc 1260
atcattgtag acatcaccaa ggacacgttt tacaaacage ccatgttcta
ccaccttggc 1320
catttcagca agttcattcc tgagggctcc cagagagtgg ggctgggtgc
cagtcagaag 1380
aacgacctgg acgcagtggc attgatgcat cccgatggct ctgctgttgt
ggtcgtgcta 1440
aaccgctcct ctaaggatgt gcctcttacc atcaaggatc ctgctgtggg
cttcctggag 1500
acaatctcac ctggctactc cattcacacc tacctgtggc gtcgccagtg a
1551

```

<210> 3 <211> 6186 <212> DNA <213> Artificial sequence  
 <220> <221> exon <222> (1225)..(1572) <223> Coding sequence  
 for human FSH-alpha <400> 3

```

gacggatcgg gagatctccc gatcccctat ggtcgactct cagtacaatc
tgctctgatg 60

```

```

ccgcatagtt aagccagtat ctgctccctg cttgtgtggt ggaggtcgct
gagtagtgcg 120

```

```

cgagcaaaat ttaagctaca acaaggcaag gcttgaccga caattgcatg
aagaatctgc 180

```





aaccgctga tcagcctcga ctgtgccttc tagttgccag ccatctgttg  
tttgccttc 1652

ccccgtgcct tccttgacct tggaagggtgc cactcccact gtcctttcct  
aataaaatga 1712

ggaaattgca tcgcattgtc tgagtaggtg tcattctatt ctgggggggtg  
gggtggggca 1772

ggacagcaag ggggaggatt gggaagacaa tagcaggcat gctggggatg  
cgggtgggctc 1832

tatggcttct gaggcggaaa gaaccagctg gggctctagg gggtatcccc  
acgcgccctg 1892

tagcggcgca ttaagcgcgg cgggtgtggt ggttacgcgc agcgtgaccg  
ctacacttgc 1952

cagcgcccta gcgcccgtc ctttcgcttt cttcccttcc tttctcgcca  
cgttcgcccgg 2012

ctttccccgt caagctctaa atcggggcat ccctttaggg ttccgattta  
gtgctttacg 2072

gcacctcgac cccaaaaaac ttgattaggg tgatgggttca cgtagtgggc  
catcgccctg 2132

atagacggtt tttcgccctt tgacgttga gtccacgttc tttaatagt  
gactcttgtt 2192

ccaaactgga acaacactca accctatctc ggtctattct tttgatttat  
aagggatatt 2252

ggggatttcg gcctattggt taaaaaatga gctgatttaa caaaaattta  
acgcgaatta 2312

attctgtgga atgtgtgtca gttagggtgt ggaaagtccc caggctcccc  
aggcaggcag 2372

aagtatgcaa agcatgcac tcaattagtc agcaaccagg tgtggaaagt  
ccccaggctc 2432

cccagcaggc agaagtatgc aaagcatgca tctcaattag tcagcaacca  
tagtcccgc 2492

cctaactccg cccatcccgc ccctaactcc gccagttcc gccattctc  
cgcccatgg 2552

ctgactaatt ttttttattt atgcagagggc cgaggccgcc tctgcctctg  
agctattcca 2612

gaagtagtga ggaggctttt ttggaggcct aggcttttgc aaaaagctcc  
cgggagcttg 2672

tatatccatt ttcggatctg atcagcacgt gatgaaaaag cctgaactca  
ccgcgacgtc 2732

tgtcgagaag tttctgatcg aaaagttcga cagcgtctcc gacctgatgc  
agctctcgga 2792

gggcgaagaa tctcgtgctt tcagcttcga ttaggagggg cgtggatatg  
tcctgcgggt 2852

aaatagctgc gccgatgggt tctacaaaga tcgttatggt tatcggcact  
ttgcatcggc 2912

cgcgctcccg attccggaag tgcttgacat tggggaattc agcgagagcc  
tgacctattg 2972

catctcccgc cgtgcacagg gtgtcacgtt gcaagacctg cctgaaaccg  
aactgcccgc 3032

tgttctgcag ccggtcgcgg aggccatgga tgcgatcgct gcggccgatc  
ttagccagac 3092

gagcgggttc ggccattcgc gaccgcaagg aatcgggtcaa tacactacat  
ggcgtgattt 3152

catatgcgcg attgctgacg cccatgtgta tcaactggcaa actgtgatgg  
acgacaccgt 3212

cagtgcgtcc gtcgcgcagg ctctcgatga gctgatgctt tgggccgagg  
actgccccga 3272

agtccggcac ctctgtcacg cggatttcgg ctccaacaat gtcctgacgg  
acaatggccg 3332

cataacagcg gtcattgact ggagcgagggc gatgttcggg gattcccaat  
acgaggtcgc 3392

caacatcttc ttctggaggc cgtgggttggc ttgtatggag cagcagacgc  
gctacttcga 3452

gcggaggcat ccggagcttg caggatcgcc gcggctccgg gcgtatatgc  
tccgcattgg 3512

tcttgaccaa ctctatcaga gcttggttga cggcaatttc gatgatgcag  
cttgggcgca 3572

gggtcgatgc gacgcaatcg tccgatccgg agccgggact gtcgggcgta  
cacaaatcgc 3632

ccgcagaagc gcggccgtct ggaccgatgg ctgtgtagaa gtactcgccg  
atagtggaaa 3692

ccgacgcccc agcactcgtc cgagggcaaaa ggaatagcac gtgctacgag  
atttcgattc 3752

caccgcccgc ttctatgaaa ggttgggctt cggaatcggt ttccgggacg  
ccggtggat 3812

gatcctccag cgcggggatc tcatgctgga gttcttcgcc caccccaact  
tgtttattgc 3872

agcttataat ggttacaaat aaagcaatag catcaciaat ttcaciaata  
aagcattttt 3932

ttcactgcat tctagttgtg gtttgtccaa actcatcaat gtatcttatt  
atgtctgtat 3992

accgtcgacc tctagctaga gcttggcgta atcatgggtca tagctgtttc  
ctgtgtgaaa 4052

ttgttatccg ctcacaattc cacacaacat acgagccgga agcataaagt  
gtaaagcctg 4112

gggtgcctaa tgagtgaagt aactcacatt aattgcgttg cgctcactgc  
ccgctttcca 4172

gtcgggaaac ctgtcgtgcc agctgcatta atgaatcggc caacgcgcgg  
ggagaggcgg 4232

tttgcgtatt gggcgctctt ccgcttcctc gctcactgac tcgctgcgct  
cggcgttcg 4292

gctgcggcga gcggtatcag ctcactcaaa ggcggtaata cggttatcca  
cagaatcagg 4352

ggataacgca ggaaagaaca tgtgagcaaa aggccagcaa aaggccagga  
accgtaaaaa 4412

ggccgcgttg ctggcgtttt tccataggct ccgccccct gacgagcatc  
acaaaaatcg 4472

acgtcaagt cagaggtggc gaaacccgac aggactataa agataaccagg  
cgtttcccc 4532

tggaagctcc ctctgtcgct ctctgttcc gaccctgccg cttaccggat  
acctgtccgc 4592

ctttctccct tcgggaagcg tggcgctttc tcaatgctca cgctgtagggt  
atctcagttc 4652

ggtgtaggtc gttcgctcca agctgggctg tgtgcacgaa cccccgttc  
agcccgaaccg 4712

ctgcgcctta tccggttaact atcgtcttga gtccaaccgc gtaagacacg  
acttatcgcc 4772

actggcagca gccactggta acaggattag cagagcgagg tatgtaggcg  
gtgctacaga 4832

gttcttgaag tgggtggccta actacggcta cactagaagg acagtatttg  
gtatctgcgc 4892

tctgctgaag ccagttacct tcggaaaaag agttggtagc tcttgatccg  
gcaaacaaac 4952

caccgctgggt agcgggtggt tttttgtttg caagcagcag attacgcgca  
gaaaaaaagg 5012

atctcaagaa gatcctttga tcttttctac ggggtctgac gctcagtggg  
acgaaaactc 5072

acgttaagggt attttgggtca tgagattatc aaaaaggatc ttcacctaga  
tccttttaaa 5132

ttaaaaaatga agtttttaaat caatctaaag tatatatgag taaacttggt  
ctgacagtta 5192

ccaatgctta atcagtgagg cacctatctc agcgatctgt ctatttcggt  
catccatagt 5252

tgcttgactc cccgtcgtgt agataactac gatacgggag ggcttaccat  
ctggccccag 5312

tgctgcaatg ataccgcgag acccacgctc accggctcca gatttatcag  
caataaacca 5372

gccagccgga agggccgagc gcagaagtgg tcctgcaact ttatccgcct  
ccatccagtc 5432

tattaattgt tgccgggaag ctagagtaag tagttcgcca gttaatagtt  
tgcgcaacgt 5492

tgttgccatt gctacaggca tcgtggtgtc acgctcgctg tttggtatgg  
cttcattcag 5552

ctccggttcc caacgatcaa ggcgagttac atgatcccc atgttgtgca  
aaaaagcgt 5612

tagctccttc ggtcctccga tcgttgctcag aagtaagttg gccgcagtgt  
tatcactcat 5672

ggttatggca gcactgcata attctcttac tgtcatgcca tccgtaagat  
gcttttctgt 5732

gactggtgag tactcaacca agtcattctg agaatagtgt atgcggcgac  
cgagttgctc 5792

ttgccggcg tcaatacggg ataataccgc gccacatagc agaactttaa  
aagtgtcat 5852

cattggaaaa cgttcttcgg ggcgaaaact ctcaaggatc ttaccgtgt  
tgagatccag 5912

ttcgatgtaa cccactcgtg cacccaactg atcttcagca tcttttactt  
tcaccagcgt 5972

ttctgggtga gcaaaaacag gaaggcaaaa tgccgcaaaa aagggaataa  
ggcgacacg 6032

gaaatgttga atactcatac tcttctttt tcaatattat tgaagcattt  
atcagggtta 6092

ttgtctcatg agcggataca tatttgaatg tatttagaaa aataaaciaa  
taggggttcc 6152

gcgcacattt ccccgaaaag tgccacctga cgtc  
6186

<210> 4 <211> 5651 <212> DNA <213> Artificial sequence  
 <220> <221> exon <222> (1231)..(1617) <223> Coding sequence  
 for human FSH-beta <400> 4

gacggatcgg gagatctccc gatccctat ggtcgactct cagtacaatc  
 tgctctgatg 60

ccgcatagtt aagccagtat ctgctccctg cttgtgtggt ggaggtcgct  
 gagtagtgcg 120

cgagcaaaat ttaagctaca acaaggcaag gcttgaccga caattgcatg  
 aagaatctgc 180

ttagggttag gcgtttttgcg ctgcttcgcg atgtacgggc cagatatacg  
 cgttgacatt 240

gattattgac tagttattaa tagtaatcaa ttacgggggc attagttcat  
 agcccatata 300

tggagttccg cgttacataa cttacggtaa atggcccgcc tggctgaccg  
 cccaacgacc 360

cccgcccatt gacgtcaata atgacgtatg ttcccatagt aacgccaata  
 gggactttcc 420

attgacgtca atgggtggac tatttacggg aaactgccc cttggcagta  
 catcaagtgt 480

atcatatgcc aagtacgccc cctattgacg tcaatgacgg taaatggccc  
 gcctggcatt 540

atgcccagta catgacctta tgggactttc ctacttggca gtacatctac  
 gtattagtca 600

tcgctattac catggtgatg cggttttggc agtacatcaa tgggcgtgga  
 tagcggtttg 660

actcacgggg atttccaagt ctccacccca ttgacgtcaa tgggagtttg  
 ttttggcacc 720

aaaatcaacg ggactttcca aaatgtcgta acaactccgc ccattgacg  
 caaatgggcg 780

gtaggcgtgt acggtgggag gtctatataa gcagagctct ctggctaact  
 agagaacca 840

ctgcttactg gcttatcgaa attaatacga ctactatag ggagacccaa  
 gctggctagc 900







gtgctacgag atttcgattc caccgcccgc ttctatgaaa gggtgggctt  
cggaatcggt 3257

ttccgggacg ccggctggat gatcctccag cgcggggatc tcatgctgga  
gttcttcgcc 3317

caccccaact tgtttattgc agcttataat gggtacaaat aaagcaatag  
catcacaat 3377

ttcacaaata aagcattttt ttcactgcat tctagttgtg gtttgtccaa  
actcatcaat 3437

gtatcttatt atgtctgtat accgtcgacc tctagctaga gcttggcgta  
atcatgggtca 3497

tagctgtttc ctgtgtgaaa ttgttatccg ctcacaattc cacacaacat  
acgagccgga 3557

agcataaagt gttaaagcctg gggcgccctaa tgagtgcgct aactcacatt  
aattgcgttg 3617

cgctcactgc ccgctttcca gtcgggaaac ctgtcgtgcc agctgcatta  
atgaatcggc 3677

caacgcgcgg ggagaggcgg tttgcgtatt gggcgctctt ccgcttcctc  
gctcactgac 3737

tcgctgcgct cggtcgttcg gctgcggcga gcggtatcag ctactcaaa  
ggcggaata 3797

cggttatcca cagaatcagg ggataacgca ggaaagaaca tgtgagcaaa  
aggccagcaa 3857

aaggccagga accgtaaaaa ggccgcgttg ctggcgtttt tccataggct  
ccgccccct 3917

gacgagcatc acaaaaatcg acgctcaagt cagaggtggc gaaacccgac  
aggactataa 3977

agataccagg cgtttcccc tggaagctcc ctctgcgct ctctgttcc  
gaccctgccg 4037

cttaccggat acctgtccgc ctttctccct tcgggaagcg tggcgctttc  
tcaatgctca 4097

cgctgtaggt atctcagttc ggtgtaggtc gttcgctcca agctgggctg  
tgtgcacgaa 4157

cccccggttc agcccgaccg ctgcgcctta tccggttaact atcgtcttga  
gtccaacccg 4217

gtaagacacg acttatcgcc actggcagca gccactggta acaggattag  
cagagcgagg 4277

tatgtaggcg gtgctacaga gttcttgaag tgggtggccta actacggcta  
cactagaagg 4337

acagtatttg gtatctgcgc tctgctgaag ccagttacct tcggaaaaag  
agttggtagc 4397

tcttgatccg gcaaacaaac caccgctggg agcggtggtt tttttgtttg  
caagcagcag 4457

attacgcgca gaaaaaaagg atctcaagaa gatcctttga tcttttctac  
ggggtctgac 4517

gctcagtgga acgaaaactc acgttaaggg attttgggtca tgagattatc  
aaaaaggatc 4577

ttcacctaga tcctttttaa ttaaaaatga agttttaaat caatctaaag  
tatatatgag 4637

taaacttggg ctgacagtta ccaatgctta atcagtgagg cacctatctc  
agcgatctgt 4697

ctatttcggt catccatagt tgccctgactc cccgtcgtgt agataactac  
gatacgggag 4757  
ggcttaccat ctggccccag tgctgcaatg ataccgcgag acccacgctc  
accggtcca 4817

gatttatcag caataaacca gccagccgga agggccgagc gcagaagtgg  
tcctgcaact 4877

ttatccgcct ccattccagtc tattaattgt tgccgggaag ctagagtaag  
tagttcgcca 4937

gttaatagtt tgcgcaacgt tggttgccatt gctacaggca tcgtgggtgc  
acgctcgtcg 4997

tttggtatgg cttcattcag ctccgggtcc caacgatcaa ggcgagttac  
atgatcccc 5057

atgttggtgca aaaaagcggg tagctccttc ggtcctccga tcgttggtcag  
aagtaagttg 5117





446310-446360

cccgcccatt gacgtcaata atgacgtatg ttcccatagt aacgccaata  
gggactttcc 420

attgacgtca atgggtggac tatttacggt aaactgcca cttggcagta  
catcaagtgt 480

atcatatgcc aagtacgccc cctattgacg tcaatgacgg taaatggccc  
gcctggcatt 540

atgccagta catgacctta tgggactttc ctacttggca gtacatctac  
gtattagtca 600

tcgctattac catggtgatg cggttttggc agtacatcaa tgggcgtgga  
tagcggtttg 660

actcaegggg atttccaagt ctccacccca ttgacgtcaa tgggagtttg  
ttttggcacc 720

aaaatcaacg ggactttcca aaatgtcgta acaactccgc cccattgacg  
caaatgggag 780

gtaggcgtgt acggtgggag gtctatataa gcagagctct ctggctaact  
agagaacca 840

ctgcttactg gcttatcgaa attaatacga ctactatag ggagacccaa  
gctggctagc 900

ttattgcggt agtttatcac agttaaattg ctaacgcagt cagtgtttct  
gacacaacag 960

tctcgaactt aagctgcagt gactctctta aggtagcctt gcagaagttg  
gtcgtgaggc 1020

actgggcagg taagtatcaa gggtacaaga caggtttaag gagaccaata  
gaaactgggc 1080

ttgtcgagac agagaagact cttgcgtttc tgataggcac ctattggtct  
tactgacatc 1140

cactttgcct ttctctccac aggtgtccac tcccagttca attacagctc  
ttaaagctt 1200

ggtagcgagc tcggatccgc cacc atg gac tac tac cgc aag tac gcc gcc  
1251

Met Asp Tyr Tyr Arg Lys Tyr Ala Ala  
1 5

1299  
 1347  
 1395  
 1443  
 1491  
 1539  
 1587  
 1639  
 1699  
 1759

atc ttc ctg gtg acc ctg agc gtg ttc ctg cac gtg ctg cac agc gcc  
1299

Ile Phe Leu Val Thr Leu Ser Val Phe Leu His Val Leu His Ser Ala  
10 15 20 25

aac atc acc gtt aac atc acc gtg gcc ccc gac gtg cag gac tgc ccc  
1347

Asn Ile Thr Val Asn Ile Thr Val Ala Pro Asp Val Gln Asp Cys Pro  
30 35 40

gag tgc acc ctg cag gag aac ccc ttc ttc agc cag ccc ggc gcc ccc  
1395

Glu Cys Thr Leu Gln Glu Asn Pro Phe Phe Ser Gln Pro Gly Ala Pro  
45 50 55

atc ctg cag tgc atg ggc tgc tgc ttc agc cgc gcc tac ccc acc ccc  
1443

Ile Leu Gln Cys Met Gly Cys Cys Phe Ser Arg Ala Tyr Pro Thr Pro  
60 65 70

ctg cgc agc aag aag acc atg ctg gtg cag aag aac gtg acc agc gag  
1491

Leu Arg Ser Lys Lys Thr Met Leu Val Gln Lys Asn Val Thr Ser Glu  
75 80 85

agc acc tgc tgc gtg gcc aag agc tac aac cgc gtg acc gtg atg ggc  
1539

Ser Thr Cys Cys Val Ala Lys Ser Tyr Asn Arg Val Thr Val Met Gly  
90 95 100 105

ggc ttc aag gtg gag aac cac acc gcc tgc cac tgc agc acc tgc tac  
1587

Gly Phe Lys Val Glu Asn His Thr Ala Cys His Cys Ser Thr Cys Tyr  
110 115 120

tac cac aag agc taatctagag ggcccgttta aaccgctga tcagcctcga  
1639

Tyr His Lys Ser  
125

ctgtgccttc tagttgccag ccattctgttg tttgcccctc ccccgctgcct  
tccttgacct 1699

tggaaggtgc cactcccact gtccttttct aataaaatga ggaaattgca  
tcgcattgtc 1759









togtgggtgtc acgctcgctcg tttgggtatgg cttcattcag ctccgggttcc  
caacgatcaa 5599

ggcgagttac atgatcccc atgttggtgca aaaaagcggg tagctccttc  
ggtcctccga 5659

togttgtcag aagtaagttg gccgcagtgt tatcactcat ggttatggca  
gcactgcata 5719

attctcttac tgtcatgccca tccgtaagat gcttttctgt gactgggtgag  
tactcaacca 5779

agtcattctg agaatagtgt atgcggcgac cgagttgctc ttgcccggcg  
tcaatacggg 5839

ataataccgc gccacatagc agaactttaa aagtgtcat cattggaaaa  
cgttcttcgg 5899

ggcgaaaact ctcaaggatc ttaccgctgt tgagatccag ttcgatgtaa  
cccactcgtg 5959

cacccaactg atcttcagca tcttttactt tcaccagcgt ttctgggtga  
gcaaaaacag 6019

gaaggcaaaa tgccgcaaaa aagggaataa gggcgacacg gaaatgttga  
atactcatac 6079

tcttcctttt tcaatattat tgaagcattt atcagggtta ttgtctcatg  
agcggatata 6139

tatttgaatg tatttagaaa aataaacaata taggggttcc gcgcacattt  
ccccgaaaag 6199

tgccacctga cgtc  
6213

<210> 8

<211> 5

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic  
peptide



<210> 11  
<211> 11  
<212> PRT  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic  
peptide

<220>  
<221> MOD\_RES  
<222> (4)  
<223> T or S

<220>  
<221> MOD\_RES  
<222> (8)  
<223> T or S

<400> 11  
Ala Asn Ile Xaa Ala Asn Ile Xaa Ala Asn Ile  
1 5 10

<210> 12  
<211> 14  
<212> PRT  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic  
peptide

<220>  
<221> MOD\_RES  
<222> (4)  
<223> T or S

<220>  
<221> MOD\_RES  
<222> (9)  
<223> T or S

<220>  
<221> MOD\_RES



<221> MOD\_RES  
<222> (4)  
<223> T or S

<220>  
<221> MOD\_RES  
<222> (7)  
<223> T or S

<220>  
<221> MOD\_RES  
<222> (10)  
<223> T or S

<400> 14  
Ala Asn His Xaa Asn Glu Xaa Asn Ala Xaa  
1 5 10

<210> 15  
<211> 7  
<212> PRT  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic  
peptide

<220>  
<221> MOD\_RES  
<222> (7)  
<223> T or S

<400> 15  
Gly Ser Pro Ile Asn Ala Xaa  
1 5

<210> 16  
<211> 13  
<212> PRT  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic  
peptide

<220>  
<221> MOD\_RES  
<222> (7)  
<223> T or S

<220>  
<221> MOD\_RES  
<222> (13)  
<223> T or S

<400> 16  
Ala Ser Pro Ile Asn Ala Xaa Ser Pro Ile Asn Ala Xaa  
1 5 10

<210> 17  
<211> 10  
<212> PRT  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic  
peptide

<220>  
<221> MOD\_RES  
<222> (4)  
<223> T or S

<220>  
<221> MOD\_RES  
<222> (7)  
<223> T or S

<220>  
<221> MOD\_RES  
<222> (10)  
<223> T or S

<400> 17  
Ala Asn Asn Xaa Asn Tyr Xaa Asn Trp Xaa  
1 5 10

<210> 18  
<211> 13  
<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic peptide

<220>

<221> MOD\_RES

<222> (5)

<223> T or S

<220>

<221> MOD\_RES

<222> (9)

<223> T or S

<220>

<221> MOD\_RES

<222> (12)

<223> T or S

<400> 18

Ala Thr Asn Ile Xaa Leu Asn Tyr Xaa Ala Asn Xaa Thr  
1 5 10

<210> 19

<211> 13

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic peptide

<220>

<221> MOD\_RES

<222> (5)

<223> T or S

<220>

<221> MOD\_RES

<222> (9)

<223> T or S

<220>

<221> MOD\_RES

<222> (13)  
<223> T or S

<400> 19  
Ala Ala Asn Ser Xaa Gly Asn Ile Xaa Ile Asn Gly Xaa  
1 5 10

<210> 20  
<211> 13  
<212> PRT  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic  
peptide

<220>  
<221> MOD\_RES  
<222> (5)  
<223> T or S

<220>  
<221> MOD\_RES  
<222> (9)  
<223> T or S

<220>  
<221> MOD\_RES  
<222> (13)  
<223> T or S

<400> 20  
Ala Val Asn Trp Xaa Ser Asn Asp Xaa Ser Asn Ser Xaa  
1 5 10

<210> 21  
<211> 13  
<212> PRT  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic  
peptide

<220>

<221> MOD\_RES

<222> (5)

<223> T or S

<220>

<221> MOD\_RES

<222> (9)

<223> T or S

<220>

<221> MOD\_RES

<222> (13)

<223> T or S

<400> 21

Ala Val Asn Trp Xaa Ser Asn Asp Xaa Ser Asn Ser Xaa

1

5

10

<210> 22

<211> 10

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic peptide

<220>

<221> MOD\_RES

<222> (4)

<223> T or S

<220>

<221> MOD\_RES

<222> (7)

<223> T or S

<220>

<221> MOD\_RES

<222> (10)

<223> T or S

<400> 22

Ala Asn Asn Xaa Asn Tyr Xaa Asn Ser Xaa

1

5

10

Sequence

<210> 23  
<211> 10  
<212> PRT  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic peptide

<400> 23  
Ala Asn Asn Thr Asn Tyr Thr Asn Trp Thr  
1 5 10

<210> 24  
<211> 15  
<212> PRT  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Linker

<400> 24  
Gly Gly Gly Gly Ser Gly Gly Gly Gly Ser Gly Gly Gly Gly Ser  
1 5 10 15

<210> 25  
<211> 35  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Primer

<400> 25  
cgcagatctg atggctggca gcctcacagg attgc  
35

<210> 26  
<211> 37  
<212> DNA  
<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Primer

<400> 26

ccggaattcc catcactggc gacgccacag gtaggtg  
37

<210> 27

<211> 35

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Primer

<400> 27

acgcgagctc gcccttgcac ccctaaaagc ttcgg  
35

<210> 28

<211> 54

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Primer

<400> 28

gcgttgacgg cagtcagagt tgacagaagg gccagccagc aaaggatagt catg  
54

<210> 29

<211> 62

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Primer

<400> 29

ctagcatgac taccctttgc tggctggccc ttctgtcaac tctgactgcc  
gtcaacgcag 60  
ct  
62

<210> 30  
<211> 48  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Primer

<400> 30  
cctgctactg ctcccagcag cagtgaaaga gtccaaagtg gcagcatg  
48

<210> 31  
<211> 56  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Primer

<400> 31  
ctagcatgct gccacttttg actctttcac tgctgctggg agcagtagca ggagct  
56

<210> 32  
<211> 21  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Primer

<400> 32  
cagctggcca tgggtaccgc g  
21

<210> 33  
<211> 4  
<212> PRT  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: N-terminal

```

peptide addition

<400> 33
Ala Asn Ile Thr
1

<210> 34
<211> 7
<212> PRT
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: N-terminal
peptide addition

<400> 34
Ala Ser Pro Ile Asn Ala Thr
1 5

<210> 35
<211> 48
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Primer

<400> 35
tgggcatcag gtgccaacat tacagcccg cctgcatcc ctaaaagc
48

<210> 36
<211> 24
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Primer

<400> 36
tttactgttt tcgtaacagt ttg
24

```

<210> 37  
 <211> 48  
 <212> DNA  
 <213> Artificial Sequence  
  
 <220>  
 <223> Description of Artificial Sequence: Primer  
  
 <400> 37  
 gcaggggcg gctgtaatgt tggcacctga tgcccacgac actgcctg  
 48

<210> 38  
 <211> 13  
 <212> PRT  
 <213> Artificial Sequence  
  
 <220>  
 <223> Description of Artificial Sequence: Synthetic  
           peptide  
  
 <220>  
 <221> MOD\_RES  
 <222> (1)..(13)  
 <223> "Xaa" represents a variable amino acid  
  
 <400> 38  
 Ala Xaa Asn Xaa Thr Xaa Asn Xaa Thr Xaa Asn Xaa Thr  
      1                  5                 10

<210> 39  
 <211> 10  
 <212> PRT  
 <213> Artificial Sequence  
  
 <220>  
 <223> Description of Artificial Sequence: Synthetic  
           peptide  
  
 <220>  
 <221> MOD\_RES  
 <222> (1)..(10)  
 <223> "Xaa" represents a variable amino acid  
  
 <400> 39

Ala Asn Xaa Thr Asn Xaa Thr Asn Xaa Thr  
 1 5 10

<210> 40  
 <211> 81  
 <212> DNA  
 <213> Artificial Sequence

<220>  
 <221> modified\_base  
 <222> (1)..(81)  
 <223> "n" represents a, t, c, g, other or unknown

<220>  
 <223> Description of Artificial Sequence: Primer

<400> 40  
 gtgtcgtggg catcaggtgc cnsaaydns achdnaayd nsachdnsaa  
 ydnsachgcc 60  
 cgccctgca tccctaaaag c  
 81

<210> 41  
 <211> 27  
 <212> DNA  
 <213> Artificial Sequence

<220>  
 <223> Description of Artificial Sequence: Primer

<400> 42  
 ggcacctgat gccacgaca ctgcctg  
 27

<210> 43  
 <211> 68  
 <212> DNA  
 <213> Artificial Sequence

<220>  
 <223> Description of Artificial Sequence: Primer

<220>  
 <221> modified\_base

<222> (1)..(68)  
<223> "nnn" is a mixture of trinucleotide codons for all  
natural amino acid residues, except proline

<400> 43  
cgtgggcatc aggtgccaac nnnachaayn nnachaaynn nachgcccgc  
ccctgcatcc 60  
ctaaaagc  
68

<210> 44  
<211> 30  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Primer

<400> 44  
gttggcacct gatgcccacg acactgctg  
30

<210> 45  
<211> 13  
<212> PRT  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic  
peptide

<220>  
<221> MOD\_RES  
<222> (4)  
<223> variable amino acid

<220>  
<221> MOD\_RES  
<222> (12)  
<223> F or L

<400> 45  
Ala Phe Asn Xaa Thr Leu Asn Lys Thr Trp Asn Xaa Thr  
1 5 10

<210> 46  
 <211> 13  
 <212> PRT  
 <213> Artificial Sequence  
  
 <220>  
 <223> Description of Artificial Sequence: Synthetic peptide

<400> 46  
 Thr Met Asn Asn Thr Trp Asn Trp Thr Trp Asn Trp Thr  
 1 5 10

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65  
66  
67  
68  
69  
70  
71  
72  
73  
74  
75  
76  
77  
78  
79  
80  
81  
82  
83  
84  
85  
86  
87  
88  
89  
90  
91  
92  
93  
94  
95  
96  
97  
98  
99  
100  
101  
102  
103  
104  
105  
106  
107  
108  
109  
110  
111  
112  
113  
114  
115  
116  
117  
118  
119  
120  
121  
122  
123  
124  
125  
126  
127  
128  
129  
130  
131  
132  
133  
134  
135  
136  
137  
138  
139  
140  
141  
142  
143  
144  
145  
146  
147  
148  
149  
150  
151  
152  
153  
154  
155  
156  
157  
158  
159  
160  
161  
162  
163  
164  
165  
166  
167  
168  
169  
170  
171  
172  
173  
174  
175  
176  
177  
178  
179  
180  
181  
182  
183  
184  
185  
186  
187  
188  
189  
190  
191  
192  
193  
194  
195  
196  
197  
198  
199  
200  
201  
202  
203  
204  
205  
206  
207  
208  
209  
210  
211  
212  
213  
214  
215  
216  
217  
218  
219  
220  
221  
222  
223  
224  
225  
226  
227  
228  
229  
230  
231  
232  
233  
234  
235  
236  
237  
238  
239  
240  
241  
242  
243  
244  
245  
246  
247  
248  
249  
250  
251  
252  
253  
254  
255  
256  
257  
258  
259  
260  
261  
262  
263  
264  
265  
266  
267  
268  
269  
270  
271  
272  
273  
274  
275  
276  
277  
278  
279  
280  
281  
282  
283  
284  
285  
286  
287  
288  
289  
290  
291  
292  
293  
294  
295  
296  
297  
298  
299  
300  
301  
302  
303  
304  
305  
306  
307  
308  
309  
310  
311  
312  
313  
314  
315  
316  
317  
318  
319  
320  
321  
322  
323  
324  
325  
326  
327  
328  
329  
330  
331  
332  
333  
334  
335  
336  
337  
338  
339  
340  
341  
342  
343  
344  
345  
346  
347  
348  
349  
350  
351  
352  
353  
354  
355  
356  
357  
358  
359  
360  
361  
362  
363  
364  
365  
366  
367  
368  
369  
370  
371  
372  
373  
374  
375  
376  
377  
378  
379  
380  
381  
382  
383  
384  
385  
386  
387  
388  
389  
390  
391  
392  
393  
394  
395  
396  
397  
398  
399  
400  
401  
402  
403  
404  
405  
406  
407  
408  
409  
410  
411  
412  
413  
414  
415  
416  
417  
418  
419  
420  
421  
422  
423  
424  
425  
426  
427  
428  
429  
430  
431  
432  
433  
434  
435  
436  
437  
438  
439  
440  
441  
442  
443  
444  
445  
446  
447  
448  
449  
450  
451  
452  
453  
454  
455  
456  
457  
458  
459  
460  
461  
462  
463  
464  
465  
466  
467  
468  
469  
470  
471  
472  
473  
474  
475  
476  
477  
478  
479  
480  
481  
482  
483  
484  
485  
486  
487  
488  
489  
490  
491  
492  
493  
494  
495  
496  
497  
498  
499  
500  
501  
502  
503  
504  
505  
506  
507  
508  
509  
510  
511  
512  
513  
514  
515  
516  
517  
518  
519  
520  
521  
522  
523  
524  
525  
526  
527  
528  
529  
530  
531  
532  
533  
534  
535  
536  
537  
538  
539  
540  
541  
542  
543  
544  
545  
546  
547  
548  
549  
550  
551  
552  
553  
554  
555  
556  
557  
558  
559  
560  
561  
562  
563  
564  
565  
566  
567  
568  
569  
570  
571  
572  
573  
574  
575  
576  
577  
578  
579  
580  
581  
582  
583  
584  
585  
586  
587  
588  
589  
590  
591  
592  
593  
594  
595  
596  
597  
598  
599  
600  
601  
602  
603  
604  
605  
606  
607  
608  
609  
610  
611  
612  
613  
614  
615  
616  
617  
618  
619  
620  
621  
622  
623  
624  
625  
626  
627  
628  
629  
630  
631  
632  
633  
634  
635  
636  
637  
638  
639  
640  
641  
642  
643  
644  
645  
646  
647  
648  
649  
650  
651  
652  
653  
654  
655  
656  
657  
658  
659  
660  
661  
662  
663  
664  
665  
666  
667  
668  
669  
670  
671  
672  
673  
674  
675  
676  
677  
678  
679  
680  
681  
682  
683  
684  
685  
686  
687  
688  
689  
690  
691  
692  
693  
694  
695  
696  
697  
698  
699  
700  
701  
702  
703  
704  
705  
706  
707  
708  
709  
710  
711  
712  
713  
714  
715  
716  
717  
718  
719  
720  
721  
722  
723  
724  
725  
726  
727  
728  
729  
730  
731  
732  
733  
734  
735  
736  
737  
738  
739  
740  
741  
742  
743  
744  
745  
746  
747  
748  
749  
750  
751  
752  
753  
754  
755  
756  
757  
758  
759  
760  
761  
762  
763  
764  
765  
766  
767  
768  
769  
770  
771  
772  
773  
774  
775  
776  
777  
778  
779  
780  
781  
782  
783  
784  
785  
786  
787  
788  
789  
790  
791  
792  
793  
794  
795  
796  
797  
798  
799  
800  
801  
802  
803  
804  
805  
806  
807  
808  
809  
810  
811  
812  
813  
814  
815  
816  
817  
818  
819  
820  
821  
822  
823  
824  
825  
826  
827  
828  
829  
830  
831  
832  
833  
834  
835  
836  
837  
838  
839  
840  
841  
842  
843  
844  
845  
846  
847  
848  
849  
850  
851  
852  
853  
854  
855  
856  
857  
858  
859  
860  
861  
862  
863  
864  
865  
866  
867  
868  
869  
870  
871  
872  
873  
874  
875  
876  
877  
878  
879  
880  
881  
882  
883  
884  
885  
886  
887  
888  
889  
890  
891  
892  
893  
894  
895  
896  
897  
898  
899  
900  
901  
902  
903  
904  
905  
906  
907  
908  
909  
910  
911  
912  
913  
914  
915  
916  
917  
918  
919  
920  
921  
922  
923  
924  
925  
926  
927  
928  
929  
930  
931  
932  
933  
934  
935  
936  
937  
938  
939  
940  
941  
942  
943  
944  
945  
946  
947  
948  
949  
950  
951  
952  
953  
954  
955  
956  
957  
958  
959  
960  
961  
962  
963  
964  
965  
966  
967  
968  
969  
970  
971  
972  
973  
974  
975  
976  
977  
978  
979  
980  
981  
982  
983  
984  
985  
986  
987  
988  
989  
990  
991  
992  
993  
994  
995  
996  
997  
998  
999  
1000  
1001  
1002  
1003  
1004  
1005  
1006  
1007  
1008  
1009  
1010  
1011  
1012  
1013  
1014  
1015  
1016  
1017  
1018  
1019  
1020  
1021  
1022  
1023  
1024  
1025  
1026  
1027  
1028  
1029  
1030  
1031  
1032  
1033  
1034  
1035  
1036  
1037  
1038  
1039  
1040  
1041  
1042  
1043  
1044  
1045  
1046  
1047  
1048  
1049  
1050  
1051  
1052  
1053  
1054  
1055  
1056  
1057  
1058  
1059  
1060  
1061  
1062  
1063  
1064  
1065  
1066  
1067  
1068  
1069  
1070  
1071  
1072  
1073  
1074  
1075  
1076  
1077  
1078  
1079  
1080  
1081  
1082  
1083  
1084  
1085  
1086  
1087  
1088  
1089  
1090  
1091  
1092  
1093  
1094  
1095  
1096  
1097  
1098  
1099  
1100  
1101  
1102  
1103  
1104  
1105  
1106  
1107  
1108  
1109  
1110  
1111  
1112  
1113  
1114  
1115  
1116  
1117  
1118  
1119  
1120  
1121  
1122  
1123  
1124  
1125  
1126  
1127  
1128  
1129  
1130  
1131  
1132  
1133  
1134  
1135  
1136  
1137  
1138  
1139  
1140  
1141  
1142  
1143  
1144  
1145  
1146  
1147  
1148  
1149  
1150  
1151  
1152  
1153  
1154  
1155  
1156  
1157  
1158  
1159  
1160  
1161  
1162  
1163  
1164  
1165  
1166  
1167  
1168  
1169  
1170  
1171  
1172  
1173  
1174  
1175  
1176  
1177  
1178  
1179  
1180  
1181  
1182  
1183  
1184  
1185  
1186  
1187  
1188  
1189  
1190  
1191  
1192  
1193  
1194  
1195  
1196  
1197  
1198  
1199  
1200  
1201  
1202  
1203  
1204  
1205  
1206  
1207  
1208  
1209  
1210  
1211  
1212  
1213  
1214  
1215  
1216  
1217  
1218  
1219  
1220  
1221  
1222  
1223  
1224  
1225  
1226  
1227  
1228  
1229  
1230  
1231  
1232  
1233  
1234  
1235  
1236  
1237  
1238  
1239  
1240  
1241  
1242  
1243  
1244  
1245  
1246  
1247  
1248  
1249  
1250  
1251  
1252  
1253  
1254  
1255  
1256  
1257  
1258  
1259  
1260  
1261  
1262  
1263  
1264  
1265  
1266  
1267  
1268  
1269  
1270  
1271  
1272  
1273  
1274  
1275  
1276  
1277  
1278  
1279  
1280  
1281  
1282  
1283  
1284  
1285  
1286  
1287  
1288  
1289  
1290  
1291  
1292  
1293  
1294  
1295  
1296  
1297  
1298  
1299  
1300  
1301  
1302  
1303  
1304  
1305  
1306  
1307  
1308  
1309  
1310  
1311  
1312  
1313  
1314  
1315  
1316  
1317  
1318  
1319  
1320  
1321  
1322  
1323  
1324  
1325  
1326  
1327  
1328  
1329  
1330  
1331  
1332  
1333  
1334  
1335  
1336  
1337  
1338  
1339  
1340  
1341  
1342  
1343  
1344  
1345  
1346  
1347  
1348  
1349  
1350  
1351  
1352  
1353  
1354  
1355  
1356  
1357  
1358  
1359  
1360  
1361  
1362  
1363  
1364  
1365  
1366  
1367  
1368  
1369  
1370  
1371  
1372  
1373  
1374  
1375  
1376  
1377  
1378  
1379  
1380  
1381  
1382  
1383  
1384  
1385  
1386  
1387  
1388  
1389  
1390  
1391  
1392  
1393  
1394  
1395  
1396  
1397  
1398  
1399  
1400  
1401  
1402  
1403  
1404  
1405  
1406  
1407  
1408  
1409  
1410  
1411  
1412  
1413  
1414  
1415  
1416  
1417  
1418  
1419  
1420  
1421  
1422  
1423  
1424  
1425  
1426  
1427  
1428  
1429  
1430  
1431  
1432  
1433  
1434  
1435  
1436  
1437  
1438  
1439  
1440  
1441  
1442  
1443  
1444  
1445  
1446  
1447  
1448  
1449  
1450  
1451  
1452  
1453  
1454  
1455  
1456  
1457  
1458  
1459  
1460  
1461  
1462  
1463  
1464  
1465  
1466  
1467  
1468  
1469  
1470  
1471  
1472  
1473  
1474  
1475  
1476  
1477  
1478  
1479  
1480  
1481  
1482  
1483  
1484  
1485  
1486  
1487  
1488  
1489  
1490  
1491  
1492  
1493  
1494  
1495  
1496  
1497  
1498  
1499  
1500  
1501  
1502  
1503  
1504  
1505  
1506  
1507  
1508  
1509  
1510  
1511  
1512  
1513  
1514  
1515  
1516  
1517  
1518  
1519  
1520  
1521  
1522  
1523  
1524  
1525  
1526  
1527  
1528  
1529  
1530  
1531  
1532  
1533  
1534  
1535  
1536  
1537  
1538  
1539  
1540  
1541  
1542  
1543  
1544  
1545  
1546  
1547  
1548  
1549  
1550  
1551  
1552  
1553  
1554  
1555  
1556  
1557  
1558  
1559  
1560  
1561  
1562  
1563  
1564  
1565  
1566  
1567  
1568  
1569  
1570  
1571  
1572  
1573  
1574  
1575  
1576  
1577  
1578  
1579  
1580  
1581  
1582  
1583  
1584  
1585  
1586  
1587  
1588  
1589  
1590  
1591  
1592  
1593  
1594  
1595  
1596  
1597  
1598  
1599  
1600  
1601  
1602  
1603  
1604  
1605  
1606  
1607  
1608  
1609  
1610  
1611  
1612  
1613  
1614  
1615  
1616  
1617  
1618  
1619  
1620  
1621  
1622  
1623  
1624  
1625  
1626  
1627  
1628  
1629  
1630  
1631  
1632  
1633  
1634  
1635  
1636  
1637  
1638  
1639  
1640  
1641  
1642  
1643  
1644  
1645  
1646  
1647  
1648  
1649  
1650  
1651  
1652  
1653  
1654  
1655  
1656  
1657  
1658  
1659  
1660  
1661  
1662  
1663  
1664  
1665  
1666  
1667  
1668  
1669  
1670  
1671  
1672  
1673  
1674  
1675  
1676  
1677  
1678  
1679  
1680  
1681  
1682  
1683  
1684  
1685  
1686  
1687  
1688  
1689  
1690  
1691  
1692  
1693  
1694  
1695  
1696  
1697  
1698  
1699  
1700  
1701  
1702  
1703  
1704  
1705  
1706  
1707  
1708  
1709  
1710  
1711  
1712  
1713  
1714  
1715  
1716  
1717  
1718  
1719  
1720  
1721  
1722  
1723  
1724  
1725  
1726  
1727  
1728  
1729  
1730  
1731  
1732  
1733  
1734  
1735  
1736  
1737  
1738  
1739  
1740  
1741  
1742  
1743  
1744  
1745  
1746  
1747  
1748  
1749  
1750  
1751  
1752  
1753  
1754  
1755  
1756  
1757  
1758  
1759  
1760  
1761  
1762  
1763  
1764  
1765  
1766  
1767  
1768  
1769  
1770  
1771  
1772  
1773  
1774  
1775  
1776  
1777  
1778  
1779  
1780  
1781  
1782  
1783  
1784  
1785  
1786  
1787  
1788  
1789  
1790  
1791  
1792  
1793  
1794  
1795  
1796  
1797  
1798  
1799  
1800  
1801  
1802  
1803  
1804  
1805  
1806  
1807  
1808  
1809  
1810  
1811  
1812  
1813  
1814  
1815  
1816  
1817  
1818  
1819  
1820  
1821  
1822  
1823  
1824  
1825  
1826  
1827  
1828  
1829  
1830  
1831  
1832  
1833  
1834  
1835  
1836  
1837  
1838  
1839  
1840  
1841  
1842  
1843  
1844  
1845  
1846  
1847  
1848  
1849  
1850  
1851  
1852  
1853  
1854  
1855  
1856  
1857  
1858  
1859  
1860  
1861  
1862  
1863  
1864  
1865  
1866  
1867  
1868  
1869  
1870  
1871  
1872  
1873  
1874  
1875  
1876  
1877  
1878  
1879  
1880  
1881  
1882  
1883  
1884  
1885  
1886  
1887  
1888  
1889  
1890  
1891  
1892  
1893  
1894  
1895  
1896  
1897  
1898  
1899  
1900  
1901  
1902  
1903  
1904  
1905  
1906  
1907  
1908  
1909  
1910  
1911  
1912  
1913  
1914  
1915  
1916  
1917  
1918  
1919  
1920  
1921  
1922  
1923  
1924  
1925  
1926  
1927  
1928  
1929  
1930  
1931  
1932  
1933  
1934  
1935  
1936  
1937  
1938  
1939  
1940  
1941  
1942  
1943  
1944  
1945  
1946  
1947  
1948  
1949  
1950  
1951  
1952  
1953  
1954  
1955  
1956  
1957  
1958  
1959  
1960  
1961  
1962  
1963  
1964  
1965  
1966  
1967  
1968  
1969  
1970  
1971  
1972  
1973  
1974  
1975  
1976  
1977  
1978  
1979  
1980  
1981  
1982  
1983  
1984  
1985  
1986  
1987  
1988  
1989  
1990  
1991  
1992  
1993  
1994  
1995  
1996  
1997  
1998  
1999  
2000  
2001  
2002  
2003  
2004  
2005  
2006  
2007  
2008  
2009  
2010  
2011  
2012  
2013  
2014  
2015  
2016  
2017  
2018  
2019  
2020  
2021  
2022  
2023  
2024  
2025  
2026  
2027  
2028  
2029  
2030  
2031  
2032  
2033  
2034  
2035  
2036  
2037  
2038  
2039  
2040  
2041  
2042  
2043  
2044  
2045  
2046  
2047  
2048  
2049  
2050  
2051  
2052  
2053  
2054  
2055  
2056  
2057  
2058  
2059  
2060  
2061  
2062  
2063  
2064  
2065  
2066  
2067  
2068  
2069  
2070  
2071  
2072  
2073  
2074  
2075  
2076  
2077  
2078  
2079  
2080  
2081  
2082  
2083  
2084  
2085  
2086  
2087  
2088  
2089  
2090  
2091  
2092  
2093  
2094  
2095  
2096  
2097  
2098  
2099  
2100  
2101  
2102  
2103  
2104  
2105  
2106  
2107  
2108  
2109  
2110  
2111  
2112  
2113  
2114  
2115  
2116  
2117  
2118  
2119  
2120  
2121  
2122  
2123  
2124  
2125  
2126  
2127  
2128  
2129  
2130  
2131  
2132  
2133  
2134  
2135  
2136  
2137  
2138  
2139  
2140  
2141  
2142  
2143  
2144  
2145  
2146  
2147  
2148  
2149  
2150  
2151  
2152  
2153  
2154  
2155  
2156  
2157  
2158  
2159  
2160  
2161  
2162  
2163  
2164  
2165  
2166  
2167  
2168  
2169  
2170  
2171  
2172  
2173  
2174  
2175  
2176  
2177  
2178  
2179  
2180  
2181  
2182  
2183  
2184  
2185  
2186  
2187  
2188  
2189  
2190  
2191  
2192  
2193  
2194  
2195  
2196  
2197  
2198  
2199  
2200  
2201  
2202  
2203  
2204  
2205  
2206  
2207

<220>  
<223> Description of Artificial Sequence: Synthetic  
peptide

<400> 49  
Thr Arg Asn Val Thr Ile Asn Cys Thr Asn Ser Thr  
1 5 10

<210> 50  
<211> 13  
<212> PRT  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic  
peptide

<400> 50  
Ala Leu Asn Trp Thr Tyr Asn Gly Thr Lys Asn Val Thr  
1 5 10

<210> 51  
<211> 13  
<212> PRT  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic  
peptide

<400> 51  
Ala Ala Asn Trp Thr Val Asn Phe Thr Gly Asn Phe Thr  
1 5 10

<210> 52  
<211> 12  
<212> PRT  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic  
peptide

<220>  
<221> MOD\_RES  
<222> (2)  
<223> variable amino acid

<220>  
<221> MOD\_RES  
<222> (4)  
<223> variable amino acid

<400> 52  
Ala Xaa Asn Xaa Thr Val Asn Ser Thr Asn Val Thr  
1 5 10

<210> 53  
<211> 13  
<212> PRT  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic  
peptide

<400> 53  
Ala Asn Asn Phe Thr Phe Asn Gly Thr Leu Asn Leu Thr  
1 5 10

<210> 54  
<211> 13  
<212> PRT  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic  
peptide

<400> 54  
Ala Gly Asn Trp Thr Ala Asn Val Thr Val Asn Val Thr  
1 5 10

<210> 55  
<211> 13  
<212> PRT  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic peptide

<400> 55  
Ala Gly Asn Ser Thr Ser Asn Val Thr Gly Asn Trp Thr  
1 5 10

<210> 56  
<211> 13  
<212> PRT  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic peptide

<400> 56  
Ala Val Asn Ser Thr Met Asn Ile His Ala Ile Pro Pro  
1 5 10

<210> 57  
<211> 13  
<212> PRT  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic peptide

<400> 57  
Ala Gly Asn Gly Thr Val Asn Gly Thr Ile Asn Gly Thr  
1 5 10

<210> 58  
<211> 13  
<212> PRT  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic peptide

<220>  
<221> MOD\_RES  
<222> (8)  
<223> variable amino acid

<400> 58  
Ala Val Asn Ser Thr Gly Asn Xaa Thr Gly Asn Trp Thr  
1 5 10

<210> 59  
<211> 12  
<212> PRT  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic  
peptide

<400> 59  
Ala Gly Asn Gly Thr Asn Gly Thr Ser Asn Leu Thr  
1 5 10

<210> 60  
<211> 13  
<212> PRT  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic  
peptide

<400> 60  
Ala Met Asn Ser Thr Lys Asn Ser Thr Leu Asn Ile Thr  
1 5 10

<210> 61  
<211> 10  
<212> PRT  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic  
peptide

<400> 61  
Ala Phe Asn Tyr Thr Ser Lys Asn Ser Thr  
1 5 10

<210> 62  
<211> 13  
<212> PRT  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic  
peptide

<400> 62  
Ala Val Asn Ala Thr Met Asn Trp Thr Ala Asn Gly Thr  
1 5 10

<210> 63  
<211> 13  
<212> PRT  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic  
peptide

<400> 63  
Ala Ser Asn Ser Thr Asn Asn Gly Thr Leu Asn Ala Thr  
1 5 10

<210> 64  
<211> 13  
<212> PRT  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic  
peptide

<400> 64  
Ala Arg Asn Lys Thr Lys Asn Phe Thr Ile Asn Leu Thr  
1 5 10

<210> 65  
<211> 12  
<212> PRT  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic peptide

<400> 65  
Ala Pro Asn Ile Thr Asn Asp Thr Val Asn Met Thr  
1 5 10

<210> 66  
<211> 13  
<212> PRT  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic peptide

<400> 66  
Ala Gln Asn Lys Thr Phe Asn Phe Thr Met Asn Cys Thr  
1 5 10

<210> 67  
<211> 13  
<212> PRT  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic peptide

<400> 67  
Ala Leu Asn Val Thr Trp Asn Cys Thr Leu Asn Leu Thr  
1 5 10

<210> 68  
<211> 10  
<212> PRT  
<213> Artificial Sequence



Ala Asn Trp Thr Asn Phe Thr Asn Trp Thr  
1 5 10

<210> 72

<211> 10

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic  
peptide

<400> 72

Pro Thr Gly Leu Ile Gly Thr Asn Phe Thr  
1 5 10

<210> 73

<211> 10

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic  
peptide

<400> 73

Ala Asn Trp Thr Asn Lys Thr Asn Phe Thr  
1 5 10

<210> 74

<211> 10

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic  
peptide

<400> 74

Ala Asn Asn Thr Asn Leu Thr Asn Ala Thr  
1 5 10

<210> 75

<211> 10  
 <212> PRT  
 <213> Artificial Sequence  
  
 <220>  
 <223> Description of Artificial Sequence: Synthetic  
           peptide  
  
 <400> 75  
 Ala Asn Tyr Thr Asn Trp Thr Asn Phe Thr  
       1                              5                              10

<210> 76  
 <211> 10  
 <212> PRT  
 <213> Artificial Sequence  
  
 <220>  
 <223> Description of Artificial Sequence: Synthetic  
           peptide  
  
 <400> 76  
 Ala Asn Thr Thr Asn Gln Thr Asn Asp Thr  
       1                              5                              10

<210> 77  
 <211> 10  
 <212> PRT  
 <213> Artificial Sequence  
  
 <220>  
 <223> Description of Artificial Sequence: Synthetic  
           peptide  
  
 <400> 77  
 Ala Asn Arg Thr Asn Trp Thr Asn Thr Thr  
       1                              5                              10

<210> 78  
 <211> 10  
 <212> PRT  
 <213> Artificial Sequence  
  
 <220>

<223> Description of Artificial Sequence: Synthetic peptide

<400> 78

Pro Thr Ala Thr Asn His Thr Asn Ser Thr  
1 5 10

<210> 79

<211> 10

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic peptide

<400> 79

Ala Asn Trp Thr Asn Gln Thr Asn Gln Thr  
1 5 10

<210> 80

<211> 10

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic peptide

<400> 80

Ala Asn Trp Thr Asn Trp Thr Asn Ala Thr  
1 5 10

<210> 81

<211> 10

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic peptide

<400> 82

Ala Asn Phe Thr Asn Lys Thr Asn Met Thr

1 5 10

<210> 83  
<211> 10  
<212> PRT  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic  
peptide

<400> 83  
Ala Asn His Thr Asn Glu Thr Asn Ala Thr  
1 5 10

Sequence

<210> 84  
<211> 10  
<212> PRT  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic  
peptide

<220>  
<221> MOD\_RES  
<222> (3)  
<223> C or W

<400> 84  
Ala Asn Xaa Thr Asn Phe Thr Asn Glu Thr  
1 5 10

<210> 85  
<211> 9  
<212> PRT  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic  
peptide

<400> 85  
Ala Asn Leu Asp Lys Leu His Lys His

1

5

<210> 86

<211> 11

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic peptide

<400> 86

Ala Asn Cys Phe Thr Asn Gln Thr Asn Phe Thr

1

5

10

<210> 87

<211> 11

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic peptide

<400> 87

Ala Asn Trp Thr Asn Trp Thr Asn Glu Trp Thr

1

5

10

<210> 88

<211> 10

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic peptide

<400> 88

Ala Asn Cys Thr Asn Trp Thr Asn Cys Thr

1

5

10

<210> 89

<211> 10

<212> PRT  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic peptide

<400> 89  
Cys His Pro Tyr Asn Trp Thr Asn Trp Thr  
1 5 10

<210> 90  
<211> 10  
<212> PRT  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic peptide

<400> 90  
Ala Asn Glu Thr Asn Tyr Thr Asn Glu Thr  
1 5 10

<210> 91  
<211> 7  
<212> PRT  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic peptide

<400> 91  
Ala Asn Trp Thr Asn Trp Thr  
1 5

<210> 92  
<211> 10  
<212> PRT  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic

peptide

<400> 92

Ala Lys Pro Tyr Lys Ser Tyr Lys Phe Tyr  
1 5 10

<210> 93

<211> 10

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic  
peptide

<400> 93

Ala Asn Ile Thr Asn Lys Thr Asn Trp Thr  
1 5 10

<210> 94

<211> 10

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic  
peptide

<400> 94

Ala Asn Trp Thr Asn Met Thr Asn Ile Thr  
1 5 10

<210> 95

<211> 10

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic  
peptide

<400> 95

Ala Asn Asn Thr Asn Arg Thr Asn Phe Thr  
1 5 10

<210> 96  
 <211> 10  
 <212> PRT  
 <213> Artificial Sequence

<220>  
 <223> Description of Artificial Sequence: Synthetic peptide

<400> 96  
 Ala Asn Trp Thr Asn Trp Thr Asn Trp Thr  
 1 5 10

<210> 97  
 <211> 11  
 <212> PRT  
 <213> Artificial Sequence

<220>  
 <223> Description of Artificial Sequence: Synthetic peptide

<400> 97  
 Ala Asn Trp Arg Thr Asn His Thr Asn Lys Thr  
 1 5 10

<210> 98  
 <211> 10  
 <212> PRT  
 <213> Artificial Sequence

<220>  
 <223> Description of Artificial Sequence: Synthetic peptide

<400> 98  
 Ala Asn Gln Thr Asn Ile Thr Asn Trp Thr  
 1 5 10

<210> 99  
 <211> 11  
 <212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic peptide

<400> 99

Ala Asn Phe Thr Asn Val Ala Thr Asn Gln Thr  
1 5 10

<210> 100

<211> 10

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic peptide

<220>

<221> MOD\_RES

<222> (1)

<223> most probable amino acid

<220>

<221> MOD\_RES

<222> (2)

<223> most probable amino acid

<220>

<221> MOD\_RES

<222> (5)

<223> variable amino acid

<220>

<221> MOD\_RES

<222> (9)

<223> most probable amino acid

<400> 100

Ala Asn Thr Thr Xaa Leu Thr Asn Lys Thr  
1 5 10

<210> 101

<211> 10

<212> PRT  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic peptide

<220>  
<221> MOD\_RES  
<222> (6)  
<223> S or C

<400> 101  
Ala Asn Lys Thr Asn Xaa Thr Asn Ile Thr  
1 5 10

<210> 102  
<211> 10  
<212> PRT  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic peptide

<220>  
<221> MOD\_RES  
<222> (9)  
<223> most probable amino acid

<400> 102  
Ala Asn Trp Thr Asn Cys Thr Asn Ile Thr  
1 5 10

<210> 103  
<211> 10  
<212> PRT  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic peptide

<220>  
<221> MOD\_RES

<222> (6)  
<223> F or L

<400> 103  
Ala Asn Trp Thr Asn Xaa Thr Asn Trp Thr  
1 5 10

<210> 104  
<211> 10  
<212> PRT  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic  
peptide

<400> 104  
Cys Gln Leu Asp Arg Ser Thr Asn Glu Thr  
1 5 10

<210> 105  
<211> 10  
<212> PRT  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic  
peptide

<400> 105  
Ala Asn Asn Thr Asn Tyr Thr Asn Trp Thr  
1 5 10

<210> 106  
<211> 10  
<212> PRT  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic  
peptide

<400> 106  
Ala Asn Asn Thr Asn Tyr Thr Asn Trp Thr

1.100-30000

1	5	10
---	---	----

<210> 107  
<211> 12  
<212> PRT  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic peptide

<400> 107  
Ala Ala Asn Asp Thr Asn Trp Thr Val Asn Cys Thr  
1 5 10

<210> 108  
<211> 13  
<212> PRT  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic peptide

<400> 108  
Ala Thr Asn Ile Thr Leu Asn Tyr Thr Ala Asn Thr Thr  
1 5 10

<210> 109  
<211> 13  
<212> PRT  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic peptide

<400> 109  
Ala Ala Asn Ser Thr Gly Asn Ile Thr Ile Asn Gly Thr  
1 5 10

<210> 110  
<211> 13

<212> PRT  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic peptide

<400> 110  
Ala Val Asn Trp Thr Ser Asn Asp Thr Ser Asn Ser Thr  
1 5 10

<210> 111  
<211> 13  
<212> PRT  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic peptide

<400> 111  
Ala Ser Pro Ile Asn Ala Thr Ser Pro Ile Asn Ala Thr  
1 5 10

<210> 112  
<211> 4  
<212> PRT  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Linker

<400> 112  
Gly Gly Gly Gly  
1

<210> 113  
<211> 4  
<212> PRT  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Linker

<400> 113  
Gly Asn Ala Thr

<210> 114  
<211> 8  
<212> PRT  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic  
peptide

<400> 114  
Asn Ser Thr Gln Asn Ala Thr Ala  
1 5

<210> 115  
<211> 14  
<212> PRT  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Synthetic  
peptide

<400> 115  
Ala Asn Leu Thr Val Arg Asn Leu Thr Arg Asn Val Thr Val  
1 5 10